



[DOI]10.12016/j.issn.2096-1456.2023.12.004

· 临床研究 ·

不同矢状骨面型患者后牙颊舌向倾斜度及 Wilson 曲线曲度的 CBCT 测量研究

姚宇，谢嘉欣，熊国平，郑雨燕，翁军权，卫晓霞

深圳市人民医院(暨南大学第二临床医学院,南方科技大学第一附属医院)口腔医学中心,广东 深圳(518020)

【摘要】 目的 研究不同矢状骨面型患者的后牙颊舌向倾斜度及 Wilson 曲线曲度,探讨不同矢状骨面型患者的后牙水平向倾斜代偿机制,为骨性错殆畸形治疗中后牙倾斜度控制提供参考依据。**方法** 本研究已通过单位伦理委员会审查批准,并获得患者知情同意。选取 90 例青少年及 90 例成人正畸治疗前的 CBCT 资料作为研究对象,两组样本中各包括矢状骨面型 I 类、II 类、III 类各 30 例。分别测量每一侧由第一前磨牙至第二磨牙的所有后牙颊舌向倾斜度、第一磨牙及第二磨牙的 Wilson 曲线曲度,比较不同矢状骨面型的青少年及成人间的差异。**结果** 与 I 类骨面型成人相比,II 类骨面型成人的上颌后牙倾斜度减小,下颌后牙倾斜度增大,差异有统计学意义;III 类骨面型成人上颌后牙倾斜度增大,下颌后牙倾斜度减小,差异有统计学意义;II 类骨面型成人第二磨牙的 Wilson 曲线曲度显著性增大,差异有统计学意义。与 I 类骨面型青少年相比,III 类骨面型青少年的上颌后牙倾斜度增大,差异有统计学意义。不同矢状骨面型青少年下颌后牙倾斜度差异无统计学意义。对比青少年与成人,II 类骨面型成人上后牙更偏舌倾,除下颌第一磨牙外下颌后牙更偏直立;III 类骨面型成人下颌后牙除下颌第二磨牙外更偏舌倾,差异有统计学意义,上颌后牙无明显变化,差异无统计学意义。**结论** 上下后牙的颊舌向倾斜度及 Wilson 曲线曲度在不同矢状骨面型人群间存在差异,II 类骨面型人群以上后牙代偿性舌倾、下后牙代偿性直立为主。III 类骨面型人群以下后牙代偿性舌倾为主,上后牙维持原有的代偿性颊倾。

【关键词】 青少年; 成人; 正畸; 后牙; 颊舌向倾斜度; Wilson 曲线曲度; 矢状骨面型; 锥形束 CT; 骨性错殆; 代偿机制



微信公众号

【中图分类号】 R78 **【文献标志码】** A **【文章编号】** 2096-1456(2023)12-0863-08

【引用著录格式】 姚宇, 谢嘉欣, 熊国平, 等. 不同矢状骨面型患者后牙颊舌向倾斜度及 Wilson 曲线曲度的 CBCT 测量研究[J]. 口腔疾病防治, 2023, 31(12): 863-870. doi:10.12016/j.issn.2096-1456.2023.12.004.

Measurement of buccolingual inclination of posterior teeth and the curve of Wilson in patients with different sagittal skeletal patterns using cone-beam computed tomography YAO Yu, XIE Jiaxin, XIONG Guoping, ZHENG Yuyan, WENG Junquan, WEI Xiaoxia. Department of Stomatology, Shenzhen People's Hospital (The Second Clinical Medical College, Jinan University; The First Affiliated Hospital, Southern University of Science and Technology), Shenzhen 518020, China

Corresponding author: WEI Xiaoxia, Email: Xiaoxiaw2007@163.com, Tel: 86-755-22943585

[Abstract] **Objective** To study the buccolingual inclination of posterior premolars and molars and the curve of Wilson in patients with different sagittal skeletal patterns, to explore the compensation mechanism of horizontal inclination of posterior teeth in patients with different sagittal skeletal patterns and to provide a reference for the control of posterior tooth inclination in the treatment of bone malocclusion. **Methods** This study was reviewed and approved by the Ethics Committee, and informed consent was obtained from the patients. Ninety CBCT scans of adults and ninety scans of adolescents before orthodontic treatment were evaluated in this cross-sectional study. There were 30 skeletal Class I,

【收稿日期】 2023-02-04; **【修回日期】** 2023-05-31

【基金项目】 国家自然科学基金项目(81900972); 深圳市科技计划项目(JCYJ20180228164315116)

【作者简介】 姚宇, 主治医师, 博士, Email: 184619127@qq.com

【通信作者】 卫晓霞, 主任医师, 博士, Email: Xiaoxiaw2007@163.com, Tel: 86-755-22943585



Class II, and Class III patients in the adult group and adolescent group. The inclination angles of posterior teeth and the curve of Wilson of first and second molars were measured, and data were analyzed between adolescents and adults with different sagittal skeletal patterns. **Results** Compared with skeletal Class I adult patients, the upper posterior molar inclination of skeletal Class II patients was significantly lower, and the lower posterior molar inclination was significantly higher. Compared with skeletal Class I adult patients, the upper posterior molar inclination of skeletal Class III adult patients was higher, and the lower posterior molar inclination was significantly lower. The Wilson curve of the second molar in skeletal Class II adult patients was significantly higher than that in the other groups. Compared with skeletal Class I adolescent patients, skeletal Class III adolescent patients had a significantly higher upper posterior molar inclination; however, no difference was found between the inclination of the posterior teeth between skeletal Class I, Class II and Class III adolescent patients. Comparing adolescent and adult samples, in skeletal Class II patients, adults showed more lingual inclination than adolescents in the upper posterior teeth and less lingual inclination in the lower posterior teeth except for the mandibular first molar. Comparing adolescent and adult samples, in skeletal Class III patients, adults showed more lingual inclination than adolescents in the lower posterior teeth except for the mandibular second molars and showed no difference in the upper posterior teeth. **Conclusions** The inclination of the posterior teeth and the curve of Wilson show significant differences between the three sagittal skeletal patterns. Compared with those of skeletal Class I patients, the posterior teeth of skeletal Class II patients show more lingual inclination in the upper arch and less lingual inclination in the lower arch. Meanwhile, posterior teeth of skeletal Class III patients show more lingual inclination in the lower arch and maintain the inclination in the upper arch.

【Key words】 adolescents; adults; orthodontics; posterior teeth; buccolingual inclination; curve of Wilson; sagittal skeletal pattern; cone-beam computer tomography; skeletal malocclusion; mechanism of compensation

J Prev Treat Stomatol Dis, 2023, 31(12): 863-870.

【Competing interests】 The authors declare no competing interests.

This study was supported by the grants from National Natural Science Foundation of China (No. 81900972); the Shenzhen Science and Technology Project (No. JCYJ20180228164315116).

牙骀结构的平衡和稳定是正畸治疗追求的目标。全口咬合的支撑依赖于正确的后牙颊舌向倾斜度。Wilson^[1]率先报道了正常天然磨牙的正常倾斜角度,即上磨牙颊倾、下磨牙舌倾。在冠状面上,倾斜的上下后牙咬合面形成一条凸向下凹向上的曲线,称之为Wilson曲线。Andrews^[2]在正常牙的六个关键中也提出,上下后牙牙冠长轴应具有一定的倾斜度,才能建立正常的后牙咬合关系。因此,充分了解后牙颊舌向倾斜度可为正畸治疗提供必要的参考。

不同的上下颌骨位置关系引导牙列在萌出建牙中发生必要代偿,不仅包括三维空间上位置的代偿,更包括倾斜度的代偿^[3-4],如Ⅱ类人群的下前牙代偿性唇倾及Ⅲ类人群的下前牙代偿性舌倾^[5]。单一维度的上下颌骨矢状向差异在三维方向上对牙骀结构产生影响。水平向上,后牙往往发生位置与倾斜度的变化^[6],但针对此问题,少有相关文献报道。因此,本研究旨在测量不同矢状骨面型青少年与成人的后牙颊舌向倾斜度,以骨性I类患者作为标准,探讨不同矢状骨面型患者

后牙在颊舌向倾斜度上的代偿机制,为正畸治疗解决上下牙弓宽度不调提供必要的参考。

1 资料和方法

本研究已获得深圳市人民医院伦理委员审批通过(批号:2022-071),已获得患者知情同意。

1.1 临床资料

选取2013年7月至2022年10月于深圳市人民医院口腔医学中心就诊的青少年(13~17岁)及成人患者(18~30岁)各90例,拍摄头颅定位侧位片及锥形束CT。根据ANB角度测量值分为骨性I类($0.7^\circ < \text{ANB} \leq 4.7^\circ$)、骨性Ⅱ类($\text{ANB} > 4.7^\circ$)、骨性Ⅲ类($\text{ANB} \leq 0.7^\circ$),各30例,共180例样本。

纳入标准:垂直骨面型为均角型,否认相关正畸治疗史;无明显后牙段拥挤不齐(拥挤度<2 mm);无后牙锁骀、反骀;无明显腭盖高拱;后牙段牙体结构健康,无明显龋坏,牙冠及牙根完整,无明显牙周组织破坏;后牙段牙齿数量正常,无缺牙(不含第三磨牙缺失)。

排除标准:上下颌骨基骨宽度差异大于5 mm;




曾有正颌外科史、颅面综合征(如唇腭裂)、面部不对称、下颌骨肥大、上呼吸道病变、上呼吸道感染、慢性口呼吸、永久性打鼾、外伤史、腺样体及扁桃体肥大等呼吸系统疾病。

1.2 测量方法

所有CBCT影像均使用NewTom VGI CBCT系统(维罗纳,意大利)扫描采集。扫描设置要求患者采用自然头位,牙尖交错位咬合。拍摄视野为 $15\text{ cm} \times 15\text{ cm}$ 。所有CT图像的空间分辨率分别为 $300\text{ }\mu\text{m}$ 、 110 kV 和 78.59 mAs 。CBCT数据使用NNT Viewer软件测量。在轴位、矢状位和冠状位对CBCT图像进行评估。在轴位及矢状位调整截面位置,在冠状位平面测量牙齿倾斜度和Wilson曲线角度。测量前重定位图像,以使眼耳平面(FH平面)和眼眶间线(连接眼眶边缘下点的线)平行于水平线。

作眼眶间线平行线,并以此作为基线,所有角度测量以此为参考。经过后牙牙冠咬合面颊舌向中心点与牙冠颈部颊舌向中心点作被测牙的牙体长轴,测量牙体长轴与基线间夹角(图1),得到以下测量值:上颌第一前磨牙倾斜度(U4I)、上颌第二前磨牙倾斜度(U5I)、上颌第一磨牙倾斜度(U6I)、上颌第二磨牙倾斜度(U7I)、下颌第一前磨牙倾斜度(L4I)、下颌第二前磨牙倾斜度(L5I)、下颌第一磨牙倾斜度(L6I)、下颌第二磨牙倾斜度(L7I)。经过上颌磨牙的近中颊尖与近中舌尖作直线,左右两侧直线相交所形成夹角角度作为Wilson曲线角度,得到第一磨牙Wilson曲线曲度(Wilson-6)和第二磨牙Wilson曲线曲度(Wilson-7)^[7]。2周后,随机抽取其中20例样本再次测量,组内相关系数最低为0.958,提示测量结果可靠,可排除测量者误差。



1.3 统计学分析

所有数据均采用SPSS18.0(SPSS公司,伊利诺伊州,美国)进行分析,数据的正态分布使用Kolmogorov-Smirnov检验进行评估。若数据呈正态分布,组间比较采用方差分析,Tukey检验用于两两比较,显著性水平为0.05。

2 结果

共采集180例样本,其中青少年90例(男52例,女38例),平均年龄(14.7 ± 2.3)岁,成人90例(男

44例,女46例),平均年龄(25.3 ± 5.2)岁。根据以往研究,矢状骨面型的性别差异并无统计学意义^[7],因此将男女数据合并统计。Kolmogorov-Smirnov检验表明所有定量变量均呈正态分布($P > 0.05$)。

2.1 不同矢状骨面型人群的后牙颊舌向倾斜度比较

90例成人患者的后牙颊舌向倾斜度见表1,不同矢状骨面型成人后牙倾斜度差异具有统计学意义。与I类骨面型成人比较,II类骨面型成人的



上颌后牙倾斜度减小,下颌后牙倾斜度增大,差异均具有统计学意义($P < 0.05$);Ⅲ类骨面型成人的上颌后牙倾斜度增大,下颌后牙倾斜度减小,差异均具有统计学意义($P < 0.05$)。不同矢状骨面型成人的第一磨牙的Wilson曲线曲度无明显差异,第二磨牙的Wilson曲线曲度差异具有统计学意义($P < 0.05$)。

表1 不同矢状骨面型成人后牙颊舌向倾斜度及Wilson曲线曲度

Table 1 The buccoligual inclination of posterior teeth and the curve of Wilson in adults with different sagittal skeletal patterns $^{\circ}, \bar{x} \pm s$

	Skeletal Clas I (n = 30)	Skeletal Clas II (n = 30)	Skeletal Class III (n = 30)	F	P
U4I	89.2 \pm 4.9 ^a	85.0 \pm 4.3 ^a	91.1 \pm 6.6 ^b	20.14	< 0.001
U5I	90.7 \pm 4.5 ^b	87.4 \pm 4.9 ^a	94.3 \pm 7.6 ^c	20.81	< 0.001
U6I	93.8 \pm 5.6 ^b	91.9 \pm 4.9 ^a	98.4 \pm 4.6 ^c	26.38	< 0.001
U7I	103.6 \pm 3.8 ^b	97.1 \pm 4.5 ^a	107.3 \pm 6.1 ^c	67.07	< 0.001
L4I	82.7 \pm 6.6 ^a	86.3 \pm 6.6 ^b	81.5 \pm 7.8 ^a	7.50	< 0.001
L5I	78.4 \pm 5.6 ^b	81.0 \pm 4.9 ^a	75.3 \pm 6.6 ^a	14.69	< 0.001
L6I	75.3 \pm 4.8 ^b	77.9 \pm 5.7 ^c	73.1 \pm 5.2 ^a	12.14	< 0.001
L7I	73.1 \pm 6.1 ^b	74.3 \pm 7.6 ^b	69.5 \pm 6.7 ^a	8.01	< 0.001
Wilson-6	157.0 \pm 8.4	160.7 \pm 8.0	157.1 \pm 8.4	1.94	0.149
Wilson-7	143.8 \pm 9.2 ^b	149.5 \pm 9.7 ^c	140.3 \pm 12.1 ^a	5.97	0.003

a < b < c was found in pairwise comparison. Significant difference was found between a and b, b and c, a and c. U4I\U5I\U6I\U7I: inclination of upper first premolar, second premolar, first molar and second molar. L4I\L5I\L6I\L7I: inclination of lower first premolar, second premolar, first molar and second molar. Wilson-6\Wilson-7: the wilson curve of first molar and second molar. Skeletal Class I: $0.7^{\circ} < ANB \leq 4.7^{\circ}$. Skeletal Class II: $ANB > 4.7^{\circ}$. Skeletal Class III: $ANB \leq 0.7^{\circ}$. The curve of Wilson: the tips of mesiobuccal and mesiolingual cusps of maxillary first and second molars along the buccal groove were connected and the formed angles were measured

90例青少年患者的后牙颊舌向倾斜度见表2,不同矢状骨面型青少年的后牙倾斜度差异亦具有统计学意义,且差异只表现在上颌,而在下颌后牙区,颊舌向倾斜度差异无统计学意义。在上颌后牙区,仅表现为Ⅲ类骨面型青少年的上颌后牙倾斜度较Ⅰ类明显增大,差异具有统计学意义($P < 0.05$);而Ⅱ类骨面型青少年的上颌后牙倾斜度与Ⅰ类差异无统计学意义。第一磨牙与第二磨牙的Wilson曲线曲度在三种矢状骨面型间的差异无统计学意义。

2.2 青少年及成人后牙颊舌向倾斜度的比较

各类矢状骨面型青少年与成人上下颌后牙的

表2 不同矢状骨面型青少年后牙颊舌向倾斜度及Wilson曲线曲度

Table 2 The buccoligual inclination of posterior teeth and the curve of Wilson in adolescents with different sagittal skeletal patterns

	skeletal patterns			$^{\circ}, \bar{x} \pm s$	
	Skeletal Clas I (n = 30)	Skeletal Clas II (n = 30)	Skeletal Class III (n = 30)	F	P
U4I	89.7 \pm 5.4 ^a	89.6 \pm 4.8 ^a	92.6 \pm 4.5 ^b	5.36	0.005
U5I	91.2 \pm 5.3 ^a	90.5 \pm 4.9 ^a	96.3 \pm 5.4 ^b	16.88	< 0.001
U6I	94.6 \pm 4.9 ^a	94.2 \pm 3.5 ^a	99.9 \pm 4.7 ^b	24.51	< 0.001
U7I	101.9 \pm 7.0 ^a	101.9 \pm 5.0 ^a	109.9 \pm 6.3 ^b	26.09	< 0.001
L4I	86.3 \pm 7.2	87.4 \pm 5.4	85.0 \pm 5.6	1.93	0.148
L5I	80.0 \pm 5.5	79.0 \pm 4.5	80.2 \pm 5.0	0.89	0.411
L6I	76.7 \pm 4.6	75.0 \pm 6.0	77.4 \pm 6.2	2.70	0.070
L7I	71.7 \pm 6.0	71.3 \pm 7.6	70.8 \pm 6.6	0.24	0.786
Wilson-6	157.0 \pm 8.4	155.0 \pm 6.5	154.7 \pm 7.8	2.40	0.099
Wilson-7	143.7 \pm 10.7	145.8 \pm 8.9	139.8 \pm 9.2	2.31	0.106

a < b < c was found in pairwise comparison. Significant difference was found between a and b, b and c, a and c. U4I\U5I\U6I\U7I: inclination of upper first premolar, second premolar, first molar and second molar. L4I\L5I\L6I\L7I: inclination of lower first premolar, second premolar, first molar and second molar. Wilson-6\Wilson-7: the wilson curve of first molar and second molar. Skeletal Class I: $0.7^{\circ} < ANB \leq 4.7^{\circ}$. Skeletal Class II: $ANB > 4.7^{\circ}$. Skeletal Class III: $ANB \leq 0.7^{\circ}$. The curve of Wilson: the tips of mesiobuccal and mesiolingual cusps of maxillary first and second molars along the buccal groove were connected and the formed angles were measured

颊舌向倾斜度见表3~表5。在Ⅰ类骨面型人群中,除下颌第一前磨牙外,成人上下颌后牙倾斜度与青少年的差异无统计学意义。

在Ⅱ类骨面型人群中,成人上颌后牙倾斜度较青少年减小,差异有统计学意义。下颌后牙除第一前磨牙外,其余后牙均表现为成人倾斜度较青少年增加,且差异具有统计学意义。成人下颌第一磨牙的Wilson曲线曲度也较青少年增大,差异具有统计学意义。第二磨牙Wilson曲线曲度在两者间的差异无统计学意义。

在Ⅲ类骨面型人群中,成人上颌后牙的颊舌向倾斜度虽然较青少年倾斜度减小,除上颌第二磨牙外,其余上颌后牙间差异均无统计学意义。在下颌后牙区,除下颌第二磨牙外的所有下颌后牙均表现为成人患者的倾斜度较青少年减小,差异具有统计学意义。成人下颌第一磨牙的Wilson曲线曲度也较青少年增大,差异具有统计学意义。第二磨牙的Wilson曲线曲度在两者间的差异无统计学意义。



表3 I类骨面型青少年与成人后牙倾斜度及Wilson曲线曲度的比较

Table 3 The comparison of buccoligual inclination of posterior teeth and the curve of Wilson in skeletal Class I

	adolescents and adults	°, $\bar{x} \pm s$		
	Adults (n = 30)	Adolescents (n = 30)	t	P
U4I	89.2 ± 4.9	89.7 ± 5.4	-0.53	0.598
U5I	90.7 ± 4.5	91.2 ± 5.3	-0.61	0.536
U6I	93.8 ± 5.6	94.6 ± 4.9	-0.83	0.407
U7I	103.6 ± 3.8	101.9 ± 7.0	1.70	0.091
L4I	82.7 ± 6.6	86.3 ± 7.2	-2.79	0.006
L5I	78.4 ± 5.6	80.0 ± 5.5	-1.56	0.121
L6I	75.3 ± 4.8	76.7 ± 4.6	-1.64	0.101
L7I	73.1 ± 6.1	71.7 ± 6.0	1.22	0.224
Wilson-6	157.7 ± 6.2	157.0 ± 8.4	0.32	0.746
Wilson-7	143.8 ± 9.2	143.7 ± 10.7	0.01	0.985

U4I\U5I\U6I\U7I: inclination of upper first premolar, second premolar, first molar and second molar. L4I\L5I\L6I\L7I: inclination of lower first premolar, second premolar, first molar and second molar. Wilson-6\Wilson-7: the wilson curve of first molar and second molar. Skeletal Class I: $0.7^\circ < ANB \leq 4.7^\circ$. Skeletal Class II: $ANB > 4.7^\circ$. Skeletal Class III: $ANB \leq 0.7^\circ$. The curve of Wilson: the tips of mesiobuccal and mesiolingual cusps of maxillary first and second molars along the buccal groove were connected and the formed angles were measured

表4 II类骨面型青少年与成人后牙倾斜度及Wilson曲线曲度的比较

Table 4 The comparison of buccoligual inclination of posterior teeth and the curve of Wilson in skeletal Class II

	adolescents and adults	°, $\bar{x} \pm s$		
	Adults (n = 30)	Adolescents (n = 30)	t	P
U4I	85.0 ± 4.3	89.6 ± 4.8	-5.43	< 0.001
U5I	87.4 ± 4.9	90.5 ± 4.9	-3.44	< 0.001
U6I	91.9 ± 4.9	94.2 ± 3.5	-3.04	0.002
U7I	97.1 ± 4.5	101.9 ± 5.0	-5.51	< 0.001
L4I	86.3 ± 6.6	87.4 ± 5.4	-1.02	0.308
L5I	81.0 ± 4.9	79.0 ± 4.5	2.33	0.021
L6I	77.9 ± 5.7	75.0 ± 6.0	2.69	0.008
L7I	74.3 ± 7.6	71.3 ± 7.6	2.19	0.029
Wilson-6	160.7 ± 8.0	155.0 ± 6.5	2.05	0.036
Wilson-7	149.5 ± 9.7	145.8 ± 8.9	1.55	0.124

U4I\U5I\U6I\U7I: inclination of upper first premolar, second premolar, first molar and second molar. L4I\L5I\L6I\L7I: inclination of lower first premolar, second premolar, first molar and second molar. Wilson-6\Wilson-7: the wilson curve of first molar and second molar. Skeletal Class I: $0.7^\circ < ANB \leq 4.7^\circ$. Skeletal Class II: $ANB > 4.7^\circ$. Skeletal Class III: $ANB \leq 0.7^\circ$. The curve of Wilson: the tips of mesiobuccal and mesiolingual cusps of maxillary first and second molars along the buccal groove were connected and the formed angles were measured

表5 III类骨面型青少年与成人后牙倾斜度及Wilson曲线曲度的比较

Table 5 The comparison of buccoligual inclination of posterior teeth and the curve of Wilson in skeletal Class III

	adolescents and adults	°, $\bar{x} \pm s$		
	Adults (n = 30)	Adolescents (n = 30)	t	P
U4I	91.1 ± 6.6	92.6 ± 4.5	-1.34	0.181
U5I	94.3 ± 7.6	96.3 ± 5.4	-1.53	0.129
U6I	98.4 ± 4.6	99.9 ± 4.7	-1.62	0.109
U7I	107.3 ± 6.1	109.9 ± 6.3	-2.02	0.047
L4I	81.5 ± 7.8	85.0 ± 5.6	-2.61	0.011
L5I	75.3 ± 6.6	80.2 ± 5.0	-4.24	< 0.001
L6I	73.1 ± 5.2	77.4 ± 6.2	-3.69	< 0.001
L7I	69.5 ± 6.7	70.8 ± 6.6	-0.95	0.344
Wilson-6	157.1 ± 8.4	154.7 ± 7.8	2.80	0.007
Wilson-7	140.3 ± 12.1	139.8 ± 9.2	0.15	0.878

U4I\U5I\U6I\U7I: inclination of upper first premolar, second premolar, first molar and second molar. L4I\L5I\L6I\L7I: inclination of lower first premolar, second premolar, first molar and second molar. Wilson-6\Wilson-7: the wilson curve of first molar and second molar. Skeletal Class I: $0.7^\circ < ANB \leq 4.7^\circ$. Skeletal Class II: $ANB > 4.7^\circ$. Skeletal Class III: $ANB \leq 0.7^\circ$. The curve of Wilson: the tips of mesiobuccal and mesiolingual cusps of maxillary first and second molars along the buccal groove were connected and the formed angles were measured

3 讨 论

影响后牙颊舌向倾斜度的因素包括年龄、矢状骨面型、垂直骨面型、基骨弓宽度、腭穹隆高度及后牙段拥挤度等^[8-10]。合理的颊舌向倾斜度是建立稳定后牙咬合的重要因素,因此区分不同骨面型及发育阶段的研究可提示后牙倾斜度在不同人群间的差异和增龄性变化下的代偿机制,为正畸治疗中协调后牙宽度不调提供参考。

锥形束CT引入颌面结构形态研究之后,基于CBCT三维重建技术的扫描测量为后牙倾斜度提供了更为精确的测量手段^[11-15]。Kasai等^[16]定义后牙牙体长轴为牙冠1/2中点及牙根尖1/3中点的连线。本研究参考Alkhatib等^[17]的研究方法,采用临床咬合面中心点及牙颈部中心点的连线作为目标牙的牙体长轴,可有效避免牙冠形态及牙根畸形变异所产生的测量误差。

既往研究中,并未发现人种差异对后牙倾斜度的影响,因此本研究未对纳入对象的人种来源作严格限定^[11-12, 17-18]。本研究纳入标准限定垂直骨面型为均角型,排除了上颌骨垂直向发育差异及腭盖深度差异,降低了腭盖深度对后牙颊舌向倾斜



度测量的干扰误差^[19]。

回顾以往研究,仅Golshah等^[7]的研究对不同矢状骨面型进行区分。其研究样本量较少,仅为66例成人,不包含青少年样本,且未对纳入样本的后牙段拥挤度作明确限制。后牙段拥挤常表现为磨牙或前磨牙的异常舌倾或颊倾,伴有锁殆、反殆,这无疑影响对代偿程度评价的准确性。因此本研究在扩大样本量及年龄层次的同时,明确规定纳入样本不存在明显后牙段拥挤。

本研究发现,不同骨面型的成人后牙颊舌向倾斜度存在差异,差异有统计学意义($P < 0.05$)。较之I类骨面型患者,I类骨面型患者的上颌后牙偏舌倾,下颌后牙偏直立。反之,III类骨面型患者的上颌后牙偏颊倾,下颌后牙偏舌倾。这提示,若上下颌骨矢状向关系协调,上下后牙可自然保持直立。若矢状向关系不调,后牙在水平向发生代偿性倾斜,以此建立咬合及最大接触面的尖窝关系。值得注意的是,II类骨面型成人的下后牙仍保留一定程度舌倾,仅较I类骨面型成人的下后牙偏直立,其倾斜度在 $86.3^\circ \sim 76.3^\circ$ 之间,尚未达到完全直立(90°),即各类矢状骨面型人群的下颌后牙均表现为舌倾,仅在舌倾幅度上存在差异。

较之I类骨面型青少年,II类骨面型青少年的后牙倾斜度均不存在明显差异,III类骨面型青少年的下后牙倾斜度同样不存在明显差异;差异最主要体现在III类骨面型青少年的上后牙,较I类明显颊倾,差异有统计学意义。这提示,无论上下颌骨矢状向是否存在不调,下颌后牙均以正常倾斜角度萌出并建殆。这可能与下颌本身的解剖生理结构有关^[20-23]。II类骨面型青少年的上颌后牙可以正常倾斜角度萌出建殆,III类骨面型青少年的上颌后牙在萌出早期即存在明显的颊倾代偿。这点在国内外研究中尚未见有相关报道。推测原因,上颌骨量较下颌更容易出现不调,II类骨面型患者上颌骨量多数正常,后牙以正常角度萌出可满足上牙列萌出需要。III类骨面型青少年伴有不同程度的上颌骨发育不足,上后牙需代偿性颊倾才能满足上牙列萌出的间隙需求。

比较青少年与成人在后牙倾斜度上的差异,发现II类骨面型成人的上后牙较青少年更偏舌倾,下后牙则更偏直立。III类骨面型的成人下后牙较青少年的更偏舌倾,上后牙无明显变化。结合上述比较结果,推测II类骨面型患者在青少年期并未表现出明显的后牙倾斜度代偿,而成人阶

段所表现的后牙倾斜可能是牙殆、骨骼、肌肉相互引导与协调,进而产生适应性代偿的结果,表现为上后牙逐步舌倾,下后牙逐步直立。III类骨面型患者则稍有不同,成人阶段的后牙仅表现为下后牙的代偿性舌倾,这可能与在青少年期就存在上颌后牙萌出方向的代偿,进而后期上后牙代偿空间不足有关。

正如Andrews正常殆六个关键理论所言,具有恰当的颊舌向倾斜度的牙冠在咬合功能中具有最大的牙尖接触面积^[2],但关于Wilson曲线在治疗后是否应该保持平坦,一直存在不同观点^[7]。因此,获得恰当的磨牙倾斜度及Wilson曲线曲度是正畸治疗中不容忽视的一部分^[24-25]。比较三种矢状骨面型人群的Wilson曲线曲度,发现第二磨牙的Wilson曲线曲度在成人间存在明显差异,II类骨面型成人的Wilson曲线最为平坦,这点与Golshah等^[7]的研究结论一致。在比较青少年的Wilson曲线曲度后,发现不同矢状骨面型人群间的差异无统计学意义。纵向比较青少年与成人的Wilson曲线曲度,结果显示I类骨面型青少年的Wilson曲线不存在明显变化,II类及III类骨面型成人第一磨牙的Wilson曲线明显较青少年平坦。这提示青少年在后牙萌出建殆初期,磨牙并未因矢状骨面型的不同而产生差异性的代偿倾斜,而在后牙长期功能性运动的影响下,磨牙的代偿性倾斜显现,Wilson曲线逐步趋向平坦。

牙殆、骨骼及肌肉等诸多因素共同参与了牙颌构建,维持着上下牙列咬合的平衡与稳定^[26-27]。正畸治疗应利于三者在三维方向上的平衡稳定,顺应生理趋势。手术病例术前正畸需去除后牙的代偿性倾斜^[28-31],匹配终末骨骼关系下的上下牙弓宽度^[32]。II类骨面型患者需颊倾上后牙、舌倾下后牙。反之,III类骨面型患者需舌倾上后牙、颊倾下后牙,这是术前正畸常见的后牙去代偿方式。针对非手术病例的掩饰治疗亦需要关注后牙倾斜度的调控。正畸减数治疗可造成牙弓不同程度缩窄,但骨性II类错殆患者的下后牙应当维持代偿性直立状态,下牙弓宽度缩窄无疑对于下颌磨牙倾斜度的控制是不利的。这也提示II类骨面型患者需注意避免因减数而造成的下颌磨牙过度舌倾,丧失后牙咬合的平衡与稳定。III类骨面型患者的掩饰性治疗,尽可能采取舌倾下后牙的策略,而非颊倾上后牙。在下颌弓丝的选择上,可以采用圆丝或小尺寸的方丝,以利于实现下后牙的代



偿性舌倾。骨性Ⅱ类青少年常伴有上颌牙弓的狭窄,但其上颌后牙并未出现明显的颊向倾斜代偿,由此推测上牙弓的缩窄可能是由于上后牙近中舌向扭转、上牙弓呈尖圆形及基骨宽度不足。在Ⅱ类骨面型青少年上颌宽度的设计中,应当以调整上颌弓形,整体扩宽上牙弓及基骨为主,避免单纯后牙颊倾形式的牙性扩弓,以期获得扩弓效果的稳定性。

由于条件限制,本研究选择的青少年与成人样本隶属于不同的样本群体,通过两类人群的结果比较推测磨牙倾斜度随年龄增长的变化趋势。在未来的研究工作中,在增加样本量的同时,可采用对青少年样本的跟踪随访,观察个体磨牙倾斜度随年龄的变化趋势,更为准确地阐述磨牙倾斜度的增龄性变化。

综上,本研究结果显示,上下后牙的颊舌向倾斜度及Wilson曲线曲度在不同矢状骨面型人群间存在差异,不同矢状骨面型人群的上下后牙在水平向上存在差异化的代偿机制。Ⅱ类骨面型人群以上后牙代偿性舌倾、下后牙代偿性直立为主。Ⅲ类骨面型人群以下后牙代偿性舌倾为主,上后牙维持原有的代偿性颊倾。这为错殆畸形的手术与非手术治疗后牙转矩控制提供了数据支持,为理解牙殆结构的三维向生理匹配与代偿机制提供了实验依据。

[Author contributions] Yao Y designed the study, processed the research, collected, analyzed the data and wrote the article. Xie JX, Xiong GP, Zheng YY, Weng JQ processed the research, collected and analyzed the data. Wei XX designed the study and reviewed the article. All authors read and approved the final manuscript as submitted.

参考文献

- [1] Wilson GH. A manual of dental prosthetics[M]. Philadelphia and New York: Lea & Febiger, 1911.
- [2] Andrews LF. The six keys to normal occlusion[J]. Am J Orthod, 1972, 62(3): 296-309. doi: 10.1016/S0002-9416(72)90268-0.
- [3] Plaza SP, Reimpell A, Silva J, et al. Relationship between skeletal Class II and Class III malocclusions with vertical skeletal pattern [J]. Dental Press J Orthod, 2019, 24(4): 63-72. doi: 10.1590/2177-6709.24.4.063-072.oar.
- [4] Chen Y, Liao W. Esthetic evaluation of incisor inclination and anterior vertical facial height[J]. Am J Orthod Dentofacial Orthop, 2022, 161(1): 3. doi: 10.1016/j.ajodo.2021.08.012.
- [5] Imamura T, Uesugi S, Ono T. Unilateral maxillary central incisor root resorption after orthodontic treatment for Angle Class II, division 1 malocclusion with significant maxillary midline deviation: a possible correlation with root proximity to the incisive canal[J]. Korean J Orthod, 2020, 50(3): 216 - 226. doi: 10.4041/kjod.2020.50.3.216.
- [6] Masumoto T, Hayashi I, Kawamura A, et al. Relationships among facial type, buccolingual molar inclination, and cortical bone thickness of the mandible[J]. Eur J Orthod, 2001, 23(1): 15-23. doi: 10.1093/ejo/23.1.15.
- [7] Golshah A, Rezaei N, Heshmati S. Buccolingual inclination of canine and first and second molar teeth and the curve of Wilson in different sagittal skeletal patterns of adults using cone-beam computed tomography[J]. Int J Dent, 2020, 2020: 8893778. doi: 10.1155/2020/8893778.
- [8] Lorente C, Lorente P, Perez-Vela M, et al. Quad-helix compression to decompensate molar inclination prior to skeletal expansion [J]. J Orofac Orthop, 2020, 81(2): 142-149. doi: 10.1007/s00056-019-00212-7.
- [9] Hwang S, Choi YJ, Jung S, et al. Posterior dental compensation and occlusal function in adults with different sagittal skeletal malocclusions[J]. Korean J Orthod, 2020, 50(2): 98-107. doi: 10.4041/kjod.2020.50.2.98.
- [10] 张婕, 马慧敏, 李小彤. 正畸-正颌联合治疗骨性Ⅲ类错殆畸形的上颌第一磨牙代偿特征初探[J]. 中华口腔医学杂志, 2018, 53(5): 307-311. doi: 10.3760/cma.j.issn.1002-0098.2018.05.004. Zhang J, Ma HM, Li XT. Compensation of upper first molar in skeletal Class Ⅲ patients treated with orthodontic-surgical treatment[J]. Zhonghua Kou Qiang Yi Xue Za Zhi, 2018, 53(5): 307-311. doi: 10.3760/cma.j.issn.1002-0098.2018.05.004.
- [11] Hu X, Huang X, Gu Y. Assessment of buccal and lingual alveolar bone thickness and buccolingual inclination of maxillary posterior teeth in patients with severe skeletal Class III malocclusion with mandibular asymmetry[J]. Am J Orthod Dentofac Orthop, 2020, 157(4): 503-515. doi: 10.1016/j.ajodo.2019.04.036.
- [12] Yang B, Chung C H. Buccolingual inclination of molars in untreated children and adults: a cone beam computed tomography study [J]. Angle Orthod, 2019, 89(1): 87-92. doi: 10.2319/010418-6.1
- [13] Moon HW, Kim MJ, Ahn HW, et al. Molar inclination and surrounding alveolar bone change relative to the design of bone-borne maxillary expanders: a CBCT study[J]. Angle Orthod, 2020, 90(1): 13-22. doi: 10.2319/050619-316.1.
- [14] Yesiltepe S, Kilci G. Evaluation the relationship between the position and impaction level of the impacted maxillary third molar teeth and marginal bone loss, caries and resorption findings of the second molar teeth with CBCT scans[J]. Oral Radiol, 2022, 38(2): 269-277. doi: 10.1007/s11282-021-00554-2.
- [15] Sharma M, Gupta S, Srivastava A, et al. CBCT analysis of maxillary first molar in Indian population[J]. Int J Clin Pediatr Dent, 2022, 15(3): 258-262. doi: 10.5005/jp-journals-10005-2375.
- [16] Kasai K, Kawamura A. Correlation between buccolingual inclination and wear of mandibular teeth in ancient and modern Japanese [J]. Arch Oral Biol, 2001, 46(3): 269 - 273. doi: 10.1016/S0003-9969(00)00106-0.
- [17] Alkhatib R, Chung CH. Buccolingual inclination of first molars in untreated adults: a CBCT study[J]. Angle Orthod, 2017, 87(4): 598-



602. doi: 10.2319/110116-786.1.
- [18] Barrera J, Llamas J, Espinar E, et al. Wilson maxillary curve analyzed by CBCT. A study on normocclusion and malocclusion individuals[J]. *Med Oral*, 2013; e547 - e552. doi: 10.4317/medoral.18291.
- [19] Prasanna AT, Saravana DS. Can palatal depth influence the buccolingual inclination of molars? A cone beam computed tomography-based retrospective evaluation[J]. *J Orthod*, 2020, 47(4): : 303-310. doi: 10.1177/1465312520941523.
- [20] Jain S, Puniyani P, Saifee A. Mandibular symphysis morphology and lower incisor angulation in different anteroposterior jaw relationships and skeletal growth patterns - a cephalometric study[J]. *Med Pharm Rep*, 2020, 93(1): 97-104. doi: 10.15386/mpr-1356.
- [21] Hu X, Huang X, Gu Y. A preliminary study of buccal and lingual alveolar bone thickness of posterior teeth in patients with skeletal Class III malocclusion and mandibular asymmetry[J]. *Am J Orthod Dentofac Orthop*, 2022, 162(1): 66 - 79. doi: 10.1016/j.ajodo.2021.02.024.
- [22] Lu CL, Li BW, Yang M, et al. Relationship between alveolar-bone morphology at the mandibular incisors and their inclination in adults with low-angle, skeletal class III malocclusion-a retrospective CBCT study[J]. *PLoS One*, 2022, 17(3): e0264788. doi:10.1371/journal.pone.0264788.
- [23] Wang J, Yang Y, Wang Y, et al. Clinical effectiveness of different types of bone-anchored maxillary protraction devices for skeletal Class III malocclusion: systematic review and network meta-analysis[J]. *Korean J Orthod*, 2022, 52(5): 313 - 323. doi:10.4041/kjod21.264.
- [24] Goh S, Dreyer C, Weir T. The predictability of the mandibular curve of Wilson, buccolingual crown inclination, and transverse expansion expression with Invisalign treatment[J]. *Am J Orthod Dentofac Orthop*, 2023, 163(1): 109 - 116. doi:10.1016/j.ajodo.2021.09.020.
- [25] Alkhafaf ZA, Sghaireen MG, Issrani R, et al. The effect of accentuation of curve of spee on masticatory efficiency-a systematic review and meta - analysis[J]. *Children (Basel)*, 2023, 10(3): 511. doi: 10.3390/children10030511.
- [26] Alkayyal MA, Turkistani KA, Al-Dharrab AA, et al. Occlusion time, occlusal balance and lateral occlusal scheme in subjects with various dental and skeletal characteristics: a prospective clinical study[J]. *J Oral Rehabil*, 2020, 47(12): 1503 - 1510. doi: 10.1111/joor.13095.
- [27] Julià-Sánchez S, Álvarez-Herms J, Cirer-Sastre R, et al. The influence of dental occlusion on dynamic balance and muscular tone [J]. *Front Physiol*, 2020, 10: 1626. doi: 10.3389/fphys.2019.01626.
- [28] Seker ED, SunalAkturk E, Yilmaz HN, et al. Occlusal plane rotation and orthodontic decompensation: influence on the outcome of surgical correction of Class III malocclusion[J]. *J Orofac Orthop Fortschritte Der Kieferorthopädie*, 2022: 1 - 11. doi: 10.1007/s00056-022-00379-6.
- [29] Quast A, Santander P, Leding J, et al. Orthodontic incisor decompensation in orthognathic therapy-success and efficiency in three dimensions[J]. *Clin Oral Invest*, 2021, 25(6): 4001 - 4010. doi: 10.1007/s00784-020-03730-6.
- [30] An JS, Jeong W, Sonnesen L, et al. Effects of presurgical mandibular incisor decompensation on long-term outcomes of class III surgical orthodontic treatment[J]. *J Clin Med*, 2021, 10(13): 2870. doi: 10.3390/jcm10132870.
- [31] Sendyk M, Cevidan LHS, de OliveiraRuellas AC, et al. Three-dimensional evaluation of dental decompensation and mandibular symphysis remodeling on orthodontic - surgical treatment of Class III malocclusion[J]. *Am J Orthod Dentofac Orthop*, 2021, 159(2): 175-183.e3. doi: 10.1016/j.ajodo.2019.12.020.
- [32] Cong A, Ruellas A, Tai SK, et al. Presurgical orthodontic decompensation with clear aligners[J]. *Am J Orthod Dentofacial Orthop*, 2022, 162(4): 538-553. doi: 10.1016/j.ajodo.2021.12.019.

(编辑 张琳,张晟)



官网