



Effect of thermal treatments on the storage stability of *Lactobacillus acidophilus* La-14 tamarind juice with or without beta-glucans

Li Yu Zhu¹, Yu Hsuan How¹, Hasanah Mohd Ghazali², Lee Sin Chang³, Kar Lin Nyam¹ and Liew Ping Pui^{1*}

¹Department of Food Science and Nutrition, Faculty of Applied Sciences, UCSI University, No. 1, Jalan Menara Gading, UCSI Heights, 56000 Cheras, Wilayah Persekutuan Kuala Lumpur, Malaysia.

²Department of Food Science, Faculty of Food Science and Technology, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia.

³Department of Food Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia.

Email: puiip@ucsiuniversity.edu.my

Received 1 March 2021; Received in revised form 20 May 2021; Accepted 6 July 2021

ABSTRACT

Aims: This study aimed to evaluate the effect of thermal treatment on the storage stability of *Lactobacillus acidophilus* La-14 tamarind juice with or without beta-glucans.

Methodology and results: *Lactobacillus acidophilus* incorporated with 6% (w/v) beta-glucans displayed the highest viability ($17.28 \log_{10}$ CFU/mL) as compared to other beta-glucans concentrations (0-8% w/v). The *L. acidophilus* with or without beta-glucans survived more than 80% after 5 h of sequential digestion. Tamarind juice was subjected to different thermal treatments (76 °C for 30 sec or 90 °C for 60 sec) and incorporated with *L. acidophilus* with or without beta-glucans. *Lactobacillus acidophilus* in tamarind juice without thermal treatment showed the highest viability ($8.69 \log_{10}$ CFU/mL), followed by thermal treatment at 76 °C for 30 sec ($>7 \log_{10}$ CFU/mL), and thermal treatment at 90 °C for 60 sec showed the lowest viability ($>4 \log_{10}$ CFU/mL), after 21 days at 4 °C. The pH, titratable acidity and viscosity of all *L. acidophilus*-tamarind juices demonstrated no changes throughout 21 days at 4 °C. Furthermore, thermal-treated tamarind juice (90 °C for 60 sec) incorporated with *L. acidophilus* displayed the least change in total soluble solids (1.99 °Brix), while thermal-treated tamarind juice (90 °C for 60 sec) with *L. acidophilus* and beta-glucans had the lowest color change ($\Delta E = 4.46$), after 21 days of storage at 4 °C.

Conclusion, significance and impact of study: Thermal treatments (90 °C for 60 sec) had contributed to the stability of *L. acidophilus*-tamarind juice with beta-glucans over 21 days of cold storage. This study shows thermal treated tamarind juice with *L. acidophilus* and beta-glucans is a potential functional non-dairy beverage catered for lactose intolerance individuals.

Keywords: Probiotic, prebiotic, functional beverage, storage stability, sequential digestion

INTRODUCTION

Probiotics are live microbiomes that can inhabit under intestinal environment and exert beneficial effect on human health (Shi *et al.*, 2016). According to Lomer *et al.* (2008), there are nearly three quarters of the total population suffers from lactose maldigestion. Owing to the fact of the high prevalence of lactose intolerance worldwide, non-dairy lactose-free probiotics product hereby plays a pronounced role in the food industry (Champagne *et al.*, 2018; Suri *et al.*, 2019).

The well-known functions of probiotics are lowering cholesterol levels, preventing antibiotic-related diarrhea and alleviating symptoms caused by lactose intolerance (Ataie-Jafari *et al.*, 2009; Fox *et al.*, 2015; Gayathri and

Vasadha, 2018). Over the past decades, researchers focused on the functionality and application of probiotics from *Lactobacillus* and *Bifidobacterium* genus (Fenster *et al.*, 2019). *Lactobacillus acidophilus* La-14 is a Gram-positive, facultative anaerobic, non-spore-forming and rod-like microorganisms (Goldstein *et al.*, 2015) that plays an important role in protecting female health (Jang *et al.*, 2017). *Lactobacillus acidophilus* La-14 possesses excellent characteristics such as high tolerance towards the harsh gastrointestinal environment, strongly adhesive to the intestinal epithelium, degrading oxalate, exert antimicrobial properties against pathogens and modulating the human immune system (Abdrabou *et al.*, 2018).

Although numerous reports had shown that probiotics

food supplements could demonstrate beneficial action in the gastrointestinal tract, combining probiotics with prebiotics can further improve their health benefits towards consumers (Mohanty *et al.*, 2018). Prebiotics are defined as non-digestible food components which can selectively facilitate the growth of one or more probiotics, therefore promoting the host's biological functions (Gibson *et al.*, 2017). It can be embedded in a wide range of foods and has rapidly drawn a lot of interest from researchers due to its high nutritional and economic value (Al-Sheraji *et al.*, 2013; Padma Ishwarya and Prabhasankar, 2014; Konar *et al.*, 2016). Beta-glucans are non-digestible dietary fiber that is found in the cell walls of microorganisms, algae and plants such as oats and barleys (Cosola *et al.*, 2017). Due to its health-promoting properties, beta-glucans were explored as a potential prebiotic to protect probiotics through harsh environments such as acid, bile and storage through functional foods (Stack *et al.*, 2010).

Emerging non-dairy probiotics food products such as probiotics nutrition bars, breakfast cereal, fruits and vegetable-based juices, and snacks have acquired a high acceptance among consumers (Granato *et al.*, 2010; Ng *et al.*, 2019; Yee *et al.*, 2019; Lai *et al.*, 2020a; Lai *et al.*, 2020b). Moreover, recent studies had suggested that fruit juices are low-fat, alcohol-free, non-dairy and nutritious beverages containing abundant natural beneficial components such as minerals, vitamins, dietary fibers and antioxidants. This allows fruit juices to be an ideal carrier for probiotics (Pereira *et al.*, 2011; Ephrem *et al.*, 2018). Tamarind pulp is rich in vitamins A, B, C and organic acids (tartaric acid, citric acid and malic acid), hence demonstrating a high antioxidant potentiality (Rai *et al.*, 2018). However, tamarind contains polyphenol oxidase enzyme that could contribute to color changes upon storage (Obulesu and Bhattacharya, 2011). Nevertheless, methods such as thermal treatment could denatured the enzymes to prevent color changes (Moon *et al.*, 2020). Furthermore, it has been reported that thermal treatment could also maintain physicochemical properties in fruit juices by inactivating undesired microorganisms (Santhirasegaram *et al.*, 2013).

Probiotics are often incorporated into dairy products that may not be suitable for lactose intolerance individuals. Hence, lactose-free fruit juices such as tamarind juice could be an alternative carrier for probiotics. In order to prevent color change and maintain physicochemical properties of the probiotic-tamarind juice during storage, different thermal treatments had been explored in this study. To date, there is yet to be any research on incorporating thermal-treated tamarind juice with probiotics and beta-glucans. Hence, this study aimed to determine the suitability of thermal-treated tamarind juice as an ideal culture medium for probiotic *L. acidophilus* growth. In addition, the prebiotic effect of beta-glucans on *L. acidophilus* La-14 through simulated gastrointestinal conditions and storage in tamarind juice was evaluated. Moreover, the physicochemical properties of the *L. acidophilus*-tamarind juice were also assessed in this study.

MATERIALS AND METHODS

Culture preparation

Lactobacillus acidophilus La-14 was prepared according to Luciano *et al.* (2018) with some modification. Briefly, 0.5 g of *L. acidophilus* La-14 powder (Pure Encapsulations, USA) was cultivated in 100 mL of sterile Man, Rogosa, Sharpe (MRS) broth (Hi-Media, India) and incubated at 37 °C for 24 h. The cells were then washed twice with phosphate-buffered saline (PBS) through centrifugation (5840 R, Eppendorf, Germany) with a speed of 4500× g for 15 min at 4 °C. The collected cell pellets were suspended in 100 mL of sterile PBS and homogenized using a vortex mixer (VTX-3000 L, LMS, Japan). The cell counts were standardized by optical density measurement at 600 nm using a spectrophotometer.

Optimization of beta-glucans concentration

Food grade beta-glucans tablets (Solgar, USA) were manually ground with a mortar and a pestle into a fine powder. The beta-glucans solution was prepared at different concentrations (0, 2, 3, 4, 5, 6, 7 and 8% w/v) by dissolving beta-glucans powder into 100 mL of MRS broth (Hi-Media, India) and autoclaved at 121 °C for 15 min (HV-110, Hirayama, Japan). After cooling down the beta-glucans mixture to ambient temperature, 1% (w/v) of *L. acidophilus* La-14 culture was inoculated into the mixture and incubated at 37 °C for 24 h (Chaikham *et al.*, 2012). The viability of *L. acidophilus* La-14 in beta-glucans solution was carried out using the drop plate method by adding 10 µL of aliquots onto MRS agar. The plates were incubated at 37 °C for 48 h and the viable cell counts of *L. acidophilus* La-14 were calculated using Equation 1 and expressed as log₁₀ CFU/mL (Phuapaiboon *et al.*, 2013). The optimal beta-glucans concentration was chosen based on the viability of *L. acidophilus* La-14 in MRS broth.

Colony forming units (CFU/mL) = Number of colonies formed / [Dilution factor × Volume plated (mL)] (1)

Preparation of simulated gastrointestinal juices

The preparation of simulated gastrointestinal juices was according to Chia *et al.* (2015) with modification. The simulated gastric juice (SGJ) was prepared by adding 3.5 mL of hydrochloric acid (Merck KGaA, Germany) and 1 g of sodium chloride (R&M Chemicals, UK) in 500 mL of distilled water. The pH of simulated gastric juice was adjusted to 2.0 using a pH meter (Eutech, USA) and sterilized at 121 °C for 15 min. After the solution was cooled down to room temperature, 1.6 g of pepsin (HmbG, Germany) was added into the mixture. On the other hand, simulated intestinal juice (SIJ) was prepared by adding 3.4 g of potassium dihydrogen phosphate (Midlothian, UK) into 95 mL of 0.1 M sodium hydroxide (Merck KGaA, Germany) and 125 mL of distilled water.

The mixture was then topped up to 500 mL with distilled water and adjusted to pH 7.5 prior to sterilizing at 121 °C for 15 min. The bile salt (R&M Chemical, UK) (3 g) was added into the mixture after it cooled down to room temperature.

Sequential digestion

Sequential digestion was carried out according to Chia *et al.* (2015) with modification. One milliliter of *L. acidophilus* La-14 with beta-glucans (6% w/v) and without beta-glucans was added into 9 mL of sterile SGJ solution, respectively. The mixture was constantly agitated at 100 rpm in the water bath (Memmert, Germany) at 37 °C for 2 h. After 2 h, the mixture was washed with PBS by centrifuging at 3200x g for 15 min at 4 °C to remove the SGJ solution. The cell pellets were then transferred to SIJ solution for another 3 h of incubation. Similarly, the mixture was also washed with PBS by centrifuging at 3200x g for 15 min at 4 °C to remove the SIJ solution. The viability of *L. acidophilus* La-14 with or without beta-glucans was carried out using the drop plate method at each hour of SGJ and SIJ incubation. Furthermore, the viable cell counts of *L. acidophilus* La-14 with or without beta-glucans at 0 h of sequential digestion was served as control. The viability was calculated using Equation 1 and expressed as log₁₀ CFU/mL. On the other hand, the survivability of *L. acidophilus* La-14 with or without beta-glucans was calculated using Equation 2 where N is the number of viable counts of cells released from microbeads at different times, and N₀ is the number of viable counts at 0 h.

$$\text{Survivability (\%)} = (\text{Log } N / \text{Log } N_0) \times 100\% \quad (2)$$

Preparation and storage stability of *L. acidophilus* La-14 in tamarind juice

The preparation of tamarind juice was in accordance with Tril *et al.* (2014) with modification. Fully ripened sour tamarind (*Tamarindus indica* LINN) was purchased from the local market. The tamarind juice was obtained by submerging 75 g of seedless tamarind pulp in 1500 mL of deionized water and blended using a juicer. The homogenized tamarind juice was then sieved three times to remove the debris. The supernatant part of the tamarind juice was collected (pH 3.6) and stored in sterile

centrifuge tubes with parafilm under 4 °C prior to thermal processing. Different thermal treatments at different temperatures (76 °C and 90 °C) and duration (30 sec and 90 sec) were used on the tamarind juice samples prior to the incorporation of 1% (w/v) *L. acidophilus* La-14 with or without 6% (w/v) beta-glucans according to Sheehan *et al.* (2007) (Table 1). All the batches of tamarind juice treated with different thermal conditions and incorporated with *L. acidophilus* with or without beta-glucans (S1, S2, S3, S4 and S5) were stored at 4 °C for 21 days.

Viability and physicochemical properties of *L. acidophilus* La-14 in tamarind juice during storage

The viability of *L. acidophilus* La-14 in tamarind juice was determined on day 0, 7, 14 and 21 of storage at 4 °C using the drop plate method with MRS agar. The agar plates were incubated at 37 °C for 48 h. The viable cell counts were calculated using Equation 1 and expressed as log₁₀ CFU/mL.

The pH, titratable acidity, total soluble solids, viscosity and color changes of *L. acidophilus* La-14 tamarind juice were evaluated on day 0, 7, 14 and 21 of storage at 4 °C. The pH was determined using a digital pH meter (pH 700, Eutech Instrument, USA). The titratable acidity was conducted according to Oladipo *et al.* (2014) where phenolphthalein was added into *L. acidophilus* La-14 tamarind juice as an indicator and titrated with 0.1 N sodium hydroxide to a pink endpoint. The titratable acidity was expressed as % lactic acid. Furthermore, the total soluble solids were measured using a refractometer (Bellingham Stanley, UK) and expressed as °Brix. Moreover, the viscosity of *L. acidophilus* La-14 tamarind juice was determined using a viscometer with spindle LV-1 at 100 rpm for 30 sec (Brookfield Viscometer DV-II+ Pro, USA) and reported in millipascal-second (mPa·s). Lastly, the total color change (ΔE) was carried out using a colorimeter (ColorFlex EZ Spectrophotometer, Hunter Colorimeter, USA) where L* (intensity of black and white), a* (intensity of red and green) and b* (intensity of yellow and blue) were read by the reflectance from the instrument. The total color change was calculated using Equation 3.

$$\text{Total color change } (\Delta E) = \sqrt{\Delta a^{*2} + \Delta b^{*2} + \Delta c^{*2}} \quad (3)$$

Table 1: Tamarind juice treated with different thermal conditions and incorporated with *L. acidophilus* La-14 with or without beta-glucans.

Tamarind juice samples	Thermal treatments		Probiotic with or without prebiotic
	Temperature (°C)	Duration (sec)	
S1	-	-	1% (w/v) <i>L. acidophilus</i> + 6% (w/v) beta-glucans
S2	76	30	1% (w/v) <i>L. acidophilus</i>
S3	76	30	1% (w/v) <i>L. acidophilus</i> + 6% (w/v) beta-glucans
S4	90	60	1% (w/v) <i>L. acidophilus</i>
S5	90	60	1% (w/v) <i>L. acidophilus</i> + 6% (w/v) beta-glucans

Statistical analysis

All data were analyzed using Minitab Statistical Analysis (version 16, Minitab Inc., USA) and reported as mean \pm standard deviation. One-way analysis of variance (ANOVA) and Tukey's post hoc test was used to evaluate the significant difference between means, where $p \leq 0.05$ was considered as significant.

RESULTS AND DISCUSSION

Optimization of beta-glucans concentration on the growth of *L. acidophilus* La-14

Figure 1 illustrates the effect of beta-glucans on the growth of *L. acidophilus* La-14. Eight different concentrations of beta-glucans were employed in this study which were 0, 2, 3, 4, 5, 6, 7 and 8% (w/v). Based on Figure 1, an increasing trend in the viability of *L. acidophilus* La-14 was observed as the concentration of beta-glucans increased. *Lactobacillus acidophilus* without the presence of beta-glucans (0% w/v) showed the lowest viability of *L. acidophilus* La-14 at 14.10 log₁₀ CFU/mL. This result is in agreement with Savedboworn *et al.* (2017) who reported that probiotic *Lactobacillus plantarum* TISTR 2075 without prebiotic (inulin) showed the lowest viability as compared to probiotics with prebiotic.

On the other hand, the viability of *L. acidophilus* La-14 increases from 15.19 to 17.28 log₁₀ CFU/mL as beta-glucans concentrations increases from 2 to 6% (w/v). However, 6 to 8% (w/v) of beta-glucans did not further increase ($p > 0.05$) the viable cell counts of *L. acidophilus* La-14. This could be due to the high concentration of beta-glucans (7-8% w/v) was not able fully utilize by the

probiotic within the 24 h incubation. Hence, the highest viability of *L. acidophilus* La-14 was reported at all 3 beta-glucans concentrations (6, 7 and 8% w/v) with the range of 17.18-17.28 log₁₀ CFU/mL. Similar results were reported by Siang *et al.* (2019) where the viable cell counts of *L. rhamnosus* GG increases as prebiotic isomalto-oligosaccharides concentration increases from 0 to 3.0% (w/v). Moreover, further addition of isomalto-oligosaccharides concentration (3 to 4% w/v) in the study also did not increase the viability of the probiotic.

In addition, the findings of this study were also in agreement with Gupta *et al.* (2017) where probiotics could acquire energy from prebiotics by consuming their end metabolic products such as acetate, propionate and butyrate. Both microbial-based beta-glucans with β -1,3, 1-6-glucans linkage and cereal-based β -1,3, 1-4-glucans can exert a positive effect on the modulation of the immune system (Rieder and Samuelsen, 2012). Besides that, beta-glucans can also form cylindrical molecules that consists approximately 0.25 million glucose units which could be utilized by probiotics as carbon source for growth (Lam and Cheung, 2013). In this study, due to similar viability ($p > 0.05$) was displayed from 6 to 8% (w/v) of beta-glucans, 6% (w/v) of beta-glucans was chosen in this optimization for further analysis due to economic reasons.

Sequential digestion for *L. acidophilus* La-14 with or without beta-glucans

The viable cell counts of *L. acidophilus* La-14 with or without beta-glucans after incubated in SGJ and SIJ for 5 h were shown in Table 2. The viability of *L. acidophilus* La-14 with or without beta-glucans had a negative correlation with the simulated sequential digestion




Figure 1: Effect of different beta-glucans concentrations on viable cell counts of *L. acidophilus* La-14. Error bars indicate the standard deviation of triplicate experiments (n=3). ^{ABCDE} means significantly different according to ANOVA and Tukey's post hoc test ($p \leq 0.05$).

Table 2: Viable cell counts and survivability of *Lactobacillus acidophilus* La-14 with or without beta-glucans under simulated gastric juice (SGJ) and simulated intestinal juice (SIJ) incubation.

Sequential digestion	Time (h)	Viable cell counts (log ₁₀ CFU/mL)		Survivability (%)	
		<i>L. acidophilus</i> La-14	<i>L. acidophilus</i> La-14 with beta-glucans	<i>L. acidophilus</i> La-14	<i>L. acidophilus</i> La-14 with beta-glucans
SGJ (pH 2)	0	14.42 ± 0.09 ^a	15.68 ± 0.06 ^a	100.00 ± 0.00 ^A	100.00 ± 0.00 ^A
	1	13.69 ± 0.03 ^b	15.65 ± 0.04 ^a	94.94 ± 0.19 ^B	99.64 ± 0.38 ^A
	2	13.67 ± 0.07 ^b	12.62 ± 0.08 ^b	94.80 ± 0.48 ^A	80.48 ± 0.43 ^B
SIJ (pH 7.5)	3	11.63 ± 0.02 ^c	12.66 ± 0.03 ^b	80.65 ± 0.11 ^A	80.74 ± 0.38 ^A
	4	11.61 ± 0.04 ^c	12.65 ± 0.03 ^b	80.51 ± 0.25 ^A	80.68 ± 0.32 ^A
	5	11.67 ± 0.04 ^c	12.60 ± 0.06 ^b	80.93 ± 0.32 ^A	80.35 ± 0.39 ^A

^{abc} Means ± standard deviation followed by different superscript letters within the same column are significantly different according to ANOVA and Tukey's post hoc test ($p \leq 0.05$). ^{AB} Means ± standard deviation followed by different superscript letters within the same row are significantly different according to ANOVA and Tukey's post hoc test ($p \leq 0.05$).

incubation time. The viable cell counts of *L. acidophilus* La-14 without beta-glucans dropped by 5.06% after the first hour of incubation in SGJ. On the other hand, no reduction in viability was found ($p > 0.05$) in *L. acidophilus* with 6% (w/v) beta-glucans after 1 h of SGJ incubation. However, during the 2nd hour of SGJ incubation, the viability of *L. acidophilus* with beta-glucans decreased by 19.4%, while *L. acidophilus* without beta-glucans remained similar viable cell counts ($p > 0.05$). This showed that beta-glucans were not effective in protecting *L. acidophilus* during the simulated gastric environment. Similar findings were reported by Yee *et al.* (2019) where prebiotic (mannitol) did not exert a prebiotic effect on probiotic (*L. acidophilus* NCFM) throughout the 2 h of SGJ incubation.

The viability of *L. acidophilus* La-14 without beta-glucans decreased by 14.9% during the first hour of SIJ incubation, while *L. acidophilus* La-14 with beta-glucans remained similar viability ($p > 0.05$). Nonetheless, the viability of both *L. acidophilus* La-14 with or without beta-glucans did not further reduce during the 2nd and 3rd hours of SIJ incubation. The reduction in viability during the first hour of SIJ incubation showed that *L. acidophilus* La-14 was sensitive to the alkaline environment and the presence of bile salt. However, *L. acidophilus* La-14 could be adapted to the stress alkaline environment with bile salt by releasing bile salt hydrolase after exposed to SIJ for 1 h (Yeung *et al.*, 2016). According to Awaisheh *et al.* (2012), exposing probiotics to stresses such as acidic, alkaline, temperature or bile salt at a milder or shorter period could enhance their survivability and tolerance towards these stresses.

Based on Table 2, *L. acidophilus* La-14 incorporated with or without 6% (w/v) beta-glucans demonstrated similar survivability of more than 80% at the end of the 5 h sequential digestion. This result demonstrated that *L. acidophilus* La-14 is a quality probiotic strain that could tolerate the environmental stress during the simulated digestive process, withstanding both stomach and intestinal pH with the presence of bile salt and digestive enzyme (Sahadeva *et al.*, 2011; Butel, 2014).

Furthermore, a standard level of viable cell counts of 10⁸–10⁹ per mL is needed to have a positive effect on modulating gastrointestinal health (Albertini *et al.*, 2010).

Both *L. acidophilus* La-14 with or without beta-glucans fulfill the required viable cell counts after sequential digestion with 11.67 and 12.60 log₁₀ CFU/mL, respectively (Table 2). This showed the acid and bile tolerance of the *L. acidophilus* La-14 strain. Although similar survivability was reported for both *L. acidophilus* La-14 with or without beta-glucans at the end of the 5 h sequential digestion, beta-glucans still showed to be an effective prebiotic in protecting *L. acidophilus* La-14 during the first hour of the simulated intestinal environment.

Viability of *L. acidophilus* La-14 with or without beta-glucans in tamarind juice during cold storage

Fruit juice is a potential vehicle for delivering probiotics due to its high nutritional value with no starter culture to compete with probiotics for growth (Nagpal *et al.*, 2012). Furthermore, oxygen scavenging components are often added into fruit juice during the manufacturing process. Therefore, processed fruit juice could often facilitate an anaerobic environment for *L. acidophilus* to grow as *L. acidophilus* is a facultatively anaerobic with optimal growth under microaerophilic conditions (Ding and Shah, 2008; Goldstein *et al.*, 2015). Table 3 shows the viability of *L. acidophilus* La-14 in different thermal-treated tamarind juice throughout 21 days of storage at 4 °C. All 5 samples of *L. acidophilus*-tamarind juice with or without beta-glucans reduced in viability throughout 21 days of cold storage.

The decreased in probiotics' viability could be due to the low pH condition of the tamarind juice (pH 3.17–3.63). According to Sheehan *et al.* (2007), the viability of probiotic (*L. paracasei*) in fruit juices is affected by pH lower than 4.5. Similar results were demonstrated by Nematollahi *et al.* (2016) and Mantzourani *et al.* (2018) where the viability of probiotics in different fruit juices reduced over cold storage. The low pH from the external

Table 3: Viability of *Lactobacillus acidophilus* La-14 tamarind juice over 21 days of storage at 4 °C.

Storage time (day)	Viable cell counts (log ₁₀ CFU/mL)				
	S1	S2	S3	S4	S5
0	14.49 ± 0.04 ^{abA}	14.21 ± 0.07 ^{abA}	14.12 ± 0.03 ^{abA}	14.18 ± 0.05 ^{abA}	14.14 ± 0.01 ^{abA}
7	11.60 ± 0.03 ^{baA}	10.86 ± 0.06 ^{bbB}	11.05 ± 0.05 ^{baA}	7.67 ± 0.10 ^{bcC}	10.02 ± 0.02 ^{bdD}
14	10.09 ± 0.08 ^{caA}	9.60 ± 0.08 ^{ccC}	10.62 ± 0.04 ^{cbB}	6.65 ± 0.02 ^{ceE}	7.81 ± 0.01 ^{cdD}
21	8.69 ± 0.08 ^{daA}	7.65 ± 0.16 ^{dbB}	7.68 ± 0.07 ^{dbB}	4.69 ± 0.11 ^{ddD}	5.68 ± 0.13 ^{dcC}

^{abcd} Means ± standard deviation followed by different superscript letters within the same column are significantly different according to ANOVA and Tukey's post hoc test ($p \leq 0.05$). ^{ABCDE} Means ± standard deviation followed by different superscript letters within the same row are significantly different according to ANOVA and Tukey's post hoc test ($p \leq 0.05$).

S1: *L. acidophilus*-tamarind juice with beta-glucans (without thermal treatment), S2: *L. acidophilus*-tamarind juice (thermal treatment 76 °C, 30 sec), S3: *L. acidophilus*-tamarind juice with beta-glucans (thermal treatment 76 °C, 30 sec), S4: *L. acidophilus*-tamarind juice (thermal treatment 90 °C, 60 sec) and S5: *L. acidophilus*-tamarind juice with beta-glucans (thermal treatment 90 °C, 60 sec).

environment could affect the survivability of probiotics by increasing the energy usage (ATP) to maintain the intracellular pH in the cells. This could further lead to insufficient ATP for other cellular activities, hence causing cell death (Nualkaekul *et al.*, 2011). Besides the pH of the fruit juice, the phytochemicals that are naturally present in tamarind juice may also exert antimicrobial activity towards the *L. acidophilus* over the storage period (Gupta *et al.*, 2014).

After 21 days of storage, S1 (tamarind juice without thermal treatment and incorporated with *L. acidophilus* La-14 and 6% w/v beta-glucans) displayed the highest viability (8.69 log₁₀ CFU/mL) and the lowest reduction of viability 40.03% as compared to the thermal-treated tamarind samples (S2, S3, S4 and S5). This could be due to the phenolic compounds and phytonutrients in the juice affected by the thermal treatments which lead to the possible reduction of nutrients availability for probiotic growth (Indrawati *et al.*, 2004; Odriozola-Serrano *et al.*, 2008; Chaikham *et al.*, 2014; Petruzzi *et al.*, 2017). Moreover, the natural antioxidant compounds of the fruit juice such as vitamin C and anthocyanins could be degraded by the thermal treatment (Chaikham *et al.*, 2014; Petruzzi *et al.*, 2017). Therefore, the increase in oxidative stress could also reduce the viability of probiotics (Odriozola-Serrano *et al.*, 2008).

The probiotic effect of beta-glucans towards *L. acidophilus* La-14 over the 21 days of cold storage was shown in Table 3. There was no difference in viability found between S2 and S3 (76 °C for 30 sec). On the other hand, *L. acidophilus* with beta-glucans in the thermal-treated tamarind juice (90 °C for 60 sec) (S5) showed higher viability (5.68 log₁₀ CFU/mL) and a lower reduction in viability (59.8%) than S4 (without beta-glucans) (4.69 log₁₀ CFU/mL) after 21 days of cold storage. This shows that beta-glucans were able to act as a nutrient source for *L. acidophilus* La-14 in thermal-treated tamarind juice (90 °C for 60 sec). The high thermal treatment could have negative effect on the nutrients in the tamarind juice (Indrawati *et al.*, 2004; Odriozola-Serrano *et al.*, 2008). Hence, the addition of probiotics can facilitate the viability of probiotics caused by acidity reduction, changes in temperature, deficient nutrients and oxidative activity (Lam and Cheung, 2013).

The final viable cell counts of *L. acidophilus* La-14 at the end of the 21 days storage for S1, S2 and S3 was more than 7 log₁₀ CFU/mL which fulfills the minimum requirement of probiotic cells in food after storage (Hill *et al.*, 2014). However, S4 and S5 did not meet the viability requirement with only 4.69 and 5.68 log₁₀ CFU/mL after 21 days of storage. This indicated that thermal treatment at 90 °C for 60 sec was not suitable for the processing of tamarind juice incorporated with *L. acidophilus*.

Physicochemical properties of *L. acidophilus* La-14 tamarind juice with or without beta-glucans during cold storage

Figure 2 shows the physicochemical changes (pH, titratable acidity, total solid content and viscosity) of *L. acidophilus* La-14 tamarind juice with or without beta-glucans during cold storage. The pH and titratable acidity of the *L. acidophilus* La-14 tamarind juice (S1, S2, S3, S4 and S5) remained constant ($p > 0.05$) throughout the 21 days of storage (Figure 2a and 2b). Fermentation is a phenomenon where probiotics could utilize sugar in fruit juice as a nutrient for growth. The organic acids produced during fermentation would decrease the pH and increase the titratable acidity of the product (Ding and Shah, 2008). However, no changes in pH or titratable acidity were observed in this study which indicates that no probiotic fermentation had occurred in the tamarind juice over 21 days of storage at 4 °C. Similar results were reported by Nematollahi *et al.* (2016) where the pH and titratable acidity of probiotic (*L. rhamnosus* or *L. plantarum*)-cornelian juice remained unchanged throughout 21 days of storage at 4 °C.

According to Ferdousi *et al.* (2013) and Savedboworn *et al.* (2019), probiotics stored under cold temperature have low cell metabolism, hence preventing the occurrence of cell activity such as the utilization of nutrients. As changes in pH and titratable acidity would affect the taste of the probiotic-fruit juice (sourness) (Mostafa *et al.*, 2021), maintaining the pH and titratable acidity throughout the storage period is crucial to provide a consistent taste of the product towards consumers. The cold temperature could offer *L. acidophilus*-tamarind juice more stable sensory properties in terms of taste.




Figure 2: Physicochemical properties of *L. acidophilus*-tamarind juice over 21 days of cold storage. (a) pH, (b) titratable acidity, (c) total soluble solids and (d) viscosity. S1: *L. acidophilus*-tamarind juice with beta-glucans (without thermal treatment), S2: *L. acidophilus*-tamarind juice (thermal treatment 76 °C, 30 sec), S3: *L. acidophilus*-tamarind juice with beta-glucans (thermal treatment 76 °C, 30 sec), S4: *L. acidophilus*-tamarind juice (thermal treatment 90 °C, 60 sec) and S5: *L. acidophilus*-tamarind juice with beta-glucans (thermal treatment 90 °C, 60 sec). Error bars indicate the standard deviation of triplicate experiments (n=3). ABCD means significantly different over 21 days of cold storage (at 4 °C) according to ANOVA and Tukey's post hoc test ($p < 0.05$).

On the other hand, total soluble solids of *L. acidophilus*-tamarind juice were reduced over 21 days of cold storage (Figure 2c). Similar results were reported by Mokhtari *et al.* (2019) and Afzaal *et al.* (2020) where total soluble solids of probiotic-fruit juice reduced over storage at 4 °C. Both studies explained that the reduction of total soluble solids was due to the utilization of sugar by probiotic during storage. However, as no reduction in pH and titratable acidity was found during the 21 days of storage in this study, therefore the reduction of total soluble solids was not due to the utilization of sugars by probiotics for growth. Non-enzymatic browning reaction is a natural chemical reaction that could occur in fruit juices upon prolonged storage. This natural phenomenon could contribute to the reduction of total soluble solids as sugar was degraded (Bharate and Bharate, 2014). Similar observation was reported by Lu *et al.* (2019) where the soluble sugar decreased upon storage.

Viscosity is one of the important parameters in evaluating beverage products as it is highly related to the

structural composition of the beverage (Damin *et al.*, 2008). The viscosity of *L. acidophilus*-tamarind juice in this study remained unchanged throughout 21 days of cold storage ($p > 0.05$) (Figure 2d). According to Garcia *et al.* (2020) and Deepak *et al.* (2016), certain lactic acid bacteria such as *L. acidophilus* could produce exopolysaccharides (EPS) that would increase the viscosity of fruit juice during fermentation. However, as pH and titratable acidity of the *L. acidophilus*-tamarind juice remained unchanged in this study, it is assumed that no fermentation had occurred. Hence, the viscosity of the *L. acidophilus*-tamarind juice remained the same throughout the cold storage. This shows that storing *L. acidophilus*-tamarind juice at 4 °C could maintain the stability of the beverage in terms of viscosity.

Based on Figure 3, the color changes in *L. acidophilus*-tamarind juice increased over 21 days of storage at 4 °C. According to Mubaiwa *et al.* (2018), the interpretation of total color changes less than 1 indicates that the color changes are not perceptible by the human




Figure 3: Total color change (ΔE) of *L. acidophilus*-tamarind juice over 21 days of cold storage at 4 °C. S1: *L. acidophilus*-tamarind juice with beta-glucans (without thermal treatment), S2: *L. acidophilus*-tamarind juice (thermal treatment 76 °C, 30 sec), S3: *L. acidophilus*-tamarind juice with beta-glucans (thermal treatment 76 °C, 30 sec), S4: *L. acidophilus*-tamarind juice (thermal treatment 90 °C, 60 sec) and S5: *L. acidophilus*-tamarind juice with beta-glucans (thermal treatment 90 °C, 60 sec).

eye, while value ranges between 1-2 means that the color changes are perceptible by the human eye via a close observation. On the other hand, total color changes locate between 2 to 10 shows that color changes can be sensed through a glance. After 7 days of storage, color change perceptible by the human eye through close observation was detected in S4 and S5 (1.92 and 1.15, respectively) which are tamarind juice that was heated at 90 °C for 60 sec, while S1, S2 and S3 had higher color change (2.17, 2.59 and 2.09, respectively) that can be sensed through a glance. The higher stability in color change displayed by tamarind juice with higher heating temperature and duration (90 °C for 60 sec) could be due to partial inactivation of enzyme present in tamarind juice that could lead to browning.

All 5 *L. acidophilus*-tamarind juice samples demonstrated color changes that can be sensed through a glance after 21 days of storage. This could be due to the breakdown of particles in tamarind juice by enzymes released by both viable and non-viable *L. acidophilus* throughout 21 days of storage (Ding and Shah, 2008; Adekunle *et al.*, 2010). Nevertheless, the least color change was displayed by S5 thermal-treated tamarind juice (90 °C for 60 sec) with the presence of *L. acidophilus* La-14 and beta-glucans. This shows that both thermal treatment and prebiotic beta-glucans could contribute to the stability of color during storage. Similar results were also reported by Renuka *et al.* (2009) where fortifying fruit juices with prebiotic fructo-oligosaccharide (FOS) were able to retain the color of the fruit juices throughout storage.

CONCLUSION

In conclusion, 6% (w/v) of beta-glucans with *L. acidophilus* La-14 displayed the highest viability (17.28 \log_{10} CFU/mL). Both *L. acidophilus* La-14 with or without beta-glucans was able to survive more than 80% after 5 h of sequential digestion. Furthermore, beta-glucans were able to protect the viability of *L. acidophilus* La-14 in tamarind juice with thermal treatment (90 °C for 60 sec) throughout 21 days of cold storage. However, thermal treatment of 76 °C for 30 sec was more suitable for tamarind juice incorporated with *L. acidophilus* La-14 than 90 °C for 60 sec as they met the minimum requirement of 6 \log_{10} CFU/mL after 21 days of storage. Cold storage (4 °C) of *L. acidophilus* La-14 tamarind juices effectively retained the pH, titratable acidity and viscosity after 21 days of storage. On the other hand, thermal treatment contributed to the stability of total soluble solids in *L. acidophilus* La-14 tamarind juices; while the combination of beta-glucans and thermal treatment was able to maintain the color stability of the *L. acidophilus* La-14 tamarind juice. Thermal-treated tamarind juice is a potential non-dairy beverage in delivering probiotic *L. acidophilus* La-14. Future studies could further explore the functionality and the sensory properties of thermal-treated tamarind juice with *L. acidophilus* La-14 and beta-glucans.

ACKNOWLEDGEMENTS

This project was supported by Pioneer Scientist Incentive Fund (PSIF) grant by UCSI University under Proj-In-Fas-

055 and UCSI Research Excellence & Innovative Grant (REIG) under REIG-FAS-2020-003.

REFERENCES

- Abdrabou, A. M., Osman, E. Y. and Aboubakr, O. A. (2018).** Comparative therapeutic efficacy study of Lactobacilli probiotics and citalopram in treatment of acute stress-induced depression in lab murine models. *Human Microbiome Journal* **10**, 33-36.
- Adekunte, A. O., Tiwari, B. K., Cullen, P. J., Scannell, A. G. M. and O'Donnell, C. P. (2010).** Effect of sonication on colour, ascorbic acid and yeast inactivation in tomato juice. *Food Chemistry* **122**(3), 500-507.
- Afzaal, M., Saeed, F., Saeed, M., Ahmed, A., Ateeq, H., Nadeem, M. T. and Tufail, T. (2020).** Survival and stability of free and encapsulated probiotic bacteria under simulated gastrointestinal conditions and in pasteurized grape juice. *Journal of Food Processing and Preservation* **44**(3), e14346.
- Al-Sheraji, S. H., Ismail, A., Manap, M. Y., Mustafa, S., Yusof, R. M. and Hassan, F. A. (2013).** Prebiotics as functional foods: A review. *Journal of Functional Foods* **5**(4), 1542-1553.
- Albertini, B., Vitali, B., Passerini, N., Cruciani, F., Di Sabatino, M., Rodriguez, L. and Brigidi, P. (2010).** Development of microparticulate systems for intestinal delivery of *Lactobacillus acidophilus* and *Bifidobacterium lactis*. *European Journal of Pharmaceutical Sciences* **40**(4), 359-366.
- Ataie-Jafari, A., Larijani, B., Alavi Majd, H. and Tahbaz, F. (2009).** Cholesterol-lowering effect of probiotic yogurt in comparison with ordinary yogurt in mildly to moderately hypercholesterolemic subjects. *Annals of Nutrition and Metabolism* **54**(1), 22-27.
- Awaisheh, S. S. (2012).** Probiotic food products classes, types, and processing. In: Probiotics. Rigobelo, E. Z. (ed.). IntechOpen, London, UK. pp. 551-582.
- Bharate, S. S. and Bharate, S. B. (2014).** Non-enzymatic browning in citrus juice: Chemical markers, their detection and ways to improve product quality. *Journal of Food Science and Technology* **51**(10), 2271-2288.
- Butel, M. J. (2014).** Probiotics, gut microbiota and health. *Médecine et Maladies Infectieuses* **44**(1), 1-8.
- Chaikhram, P., Apichartsrangkoon, A., Jirattananang, W. and Van de Wiele, T. (2012).** Influence of encapsulated probiotics combined with pressurized longan juice on colon microflora and their metabolic activities on the exposure to simulated dynamic gastrointestinal tract. *Food Research International* **49**(1), 133-142.
- Chaikhram, P., Worametrachanon, S. and Apichartsrangkoon, A. (2014).** Effects of high pressure and thermal processing on phytochemical, color and microbiological qualities of herbal-plant infusion. *International Food Research Journal* **21**(1), 51-57.
- Champagne, C. P., Gomes da Cruz, A. and Daga, M. (2018).** Strategies to improve the functionality of probiotics in supplements and foods. *Current Opinion in Food Science* **22**, 160-166.
- Chia, P. X., Tan, L. J., Huang, C. M. Y., Chan, E. W. C. and Wong, S. Y. W. (2015).** Hydrogel beads from sugar cane bagasse and palm kernel cake, and the viability of encapsulated *Lactobacillus acidophilus*. *e-Polymers* **15**(6), 411-418.
- Cosola, C., De Angelis, M., Rocchetti, M. T., Montemurno, E., Maranzano, V., Dalfino, G., Manno, C., Zito, A., Gesualdo, M., Ciccone, M. M., Gobbetti, M. and Gesualdo, L. (2017).** Beta-glucans supplementation associates with reduction in p-cresyl sulfate levels and improved endothelial vascular reactivity in healthy individuals. *PLoS ONE* **12**(1), e0169635.
- Damin, M. R., Minowa, E., Alcântara, M. R. and Oliveira, M. N. (2008).** Effect of cold storage on culture viability and some rheological properties of fermented milk prepared with yogurt and probiotic bacteria. *Journal of Texture Studies* **39**(1), 40-55.
- Deepak, V., Ram Kumar Pandian, S., Sivasubramaniam, S. D., Nellaiah, H. and Sundar, K. (2016).** Optimization of anticancer exopolysaccharide production from probiotic *Lactobacillus acidophilus* by response surface methodology. *Preparative Biochemistry and Biotechnology* **46**(3), 288-297.
- Ding, W. K. and Shah, N. P. (2008).** Survival of free and microencapsulated probiotic bacteria in orange and apple juices. *International Food Research Journal* **15**(2), 219-232.
- Ephrem, E., Najjar, A., Charcosset, C. and Greige-Gerges, H. (2018).** Encapsulation of natural active compounds, enzymes, and probiotics for fruit juice fortification, preservation, and processing: An overview. *Journal of Functional Foods* **48**, 65-84.
- Fenster, K., Freeburg, B., Hollard, C., Wong, C., Laursen, R. R. and Ouweland, A. C. (2019).** The production and delivery of probiotics: A review of a practical approach. *Microorganisms* **7**(3), 83.
- Ferdousi, R., Rouhi, M., Mohammadi, R., Mortazavian, A. M., Khosravi-Darani, K. and Rad, A. H. (2013).** Evaluation of probiotic survivability in yogurt exposed to cold chain interruption. *Iranian Journal of Pharmaceutical Research* **12**, 139-144.
- Fox, M. J., Ahuja, K. D. K., Robertson, I. K., Ball, M. J. and Eri, R. D. (2015).** Can probiotic yogurt prevent diarrhoea in children on antibiotics? A double-blind, randomised, placebo-controlled study. *BMJ Open* **5**, e006474.
- Garcia, C., Guerin, M., Souidi, S. and Remize, F. (2020).** Lactic fermented fruit or vegetable juices: Past, present and future. *Beverages* **6**(1), 8.
- Gayathri, D. and Vasadha, M. (2018).** Lactose intolerance with special emphasis on probiotics for management. *EC Nutrition* **13**(5), 325-332.
- Gibson, G. R., Hutkins, R., Sanders, M. E., Prescott, S. L., Reimer, R. A., Salminen, S. J., Scott, K., Stanton, C., Swanson, K. S., Cani, P. D., Verbeke, K. and Reid, G. (2017).** The International Scientific

- Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nature Reviews Gastroenterology and Hepatology* **14**, 491-502.
- Goldstein, E. J. C., Tyrrell, K. L. and Citron, D. M. (2015).** *Lactobacillus* species: Taxonomic complexity and controversial susceptibilities. *Clinical Infectious Diseases* **60**(Suppl 2), S98-S107.
- Granato, D., Branco, G. F., Nazzaro, F., Cruz, A. G. and Faria, J. A. F. (2010).** Functional foods and nondairy probiotic food development: Trends, concepts, and products. *Comprehensive Reviews in Food Science and Food Safety* **9**(3), 292-302.
- Gupta, C., Prakash, D. and Gupta, S. (2014).** Studies on the antimicrobial activity of tamarind (*Tamarindus indica*) and its potential as food bio-preservative. *International Food Research Journal* **21**(6), 2437-2441.
- Gupta, V. K., Paul, S. and Dutta, C. (2017).** Geography, ethnicity or subsistence-specific variations in human microbiome composition and diversity. *Frontiers in Microbiology* **8**, 1162.
- Hill, C., Guarner, F., Reid, G., Gibson, G. R., Merenstein, D. J., Pot, B., Morelli, L., Canani, R. B., Flint, H. J., Salminen, S., Calder, P. C. and Sanders, M. E. (2014).** The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nature Reviews Gastroenterology and Hepatology* **11**, 506-514.
- Indrawati, Van Loey, A. and Hendrickx, M. (2004).** Pressure and temperature stability of water-soluble antioxidants in orange and carrot juice: A kinetic study. *European Food Research and Technology* **219**(2), 161-166.
- Jang, S. E., Jeong, J. J., Choi, S. Y., Kim, H., Han, M. J. and Kim, D. H. (2017).** *Lactobacillus rhamnosus* HN001 and *Lactobacillus acidophilus* La-14 attenuate *Gardnerella vaginalis*-infected bacterial vaginosis in mice. *Nutrients* **9**(6), 531.
- Konar, N., Toker, O. S., Oba, S. and Sagdic, O. (2016).** Improving functionality of chocolate: A review on probiotic, prebiotic, and/or synbiotic characteristics. *Trends in Food Science and Technology* **49**, 35-44.
- Lai, J. T., Lai, K. W., Zhu, L. Y., Nyam, K. L. and Pui, L. P. (2020a).** Microencapsulation of *Lactobacillus plantarum* 299v and its storage in kuini juice. *Malaysian Journal of Microbiology* **16**(4), 235-244.
- Lai, K. W., How, Y. H. and Pui, L. P. (2020b).** Storage stability of microencapsulated *Lactobacillus rhamnosus* GG in hawthorn berry tea with flaxseed mucilage. *Journal of Food Processing and Preservation* **44**(12), e14965.
- Lam, K. L. and Cheung, P. C. K. (2013).** Non-digestible long chain beta-glucans as novel prebiotics. *Bioactive Carbohydrates and Dietary Fibre* **2**(1), 45-64.
- Lomer, M. C. E., Parkes, G. C. and Sanderson, J. D. (2008).** Review article: Lactose intolerance in clinical practice - Myths and realities. *Alimentary Pharmacology and Therapeutics* **27**(2), 93-103.
- Lu, Q., Li, L., Xue, S., Yang, D. and Wang, S. (2019).** Stability of flavonoid, carotenoid, soluble sugar and Vitamin C in 'Cara Cara' juice during storage. *Foods* **8**(9), 417.
- Luciano, W. A., Matte, T. C., Portela, I. A., de Medeiros, L. L., dos Santos Lima, M., Maciel, J. F., de Souza, E. L., Garcia, E. F. and Magnani, M. (2018).** Effects of *Lactobacillus acidophilus* LA-3 on physicochemical and sensory parameters of açai and mango based smoothies and its survival following simulated gastrointestinal conditions. *Food Research International* **114**, 159-168.
- Mantzourani, I., Nouska, C., Terpou, A., Alexopoulos, A., Bezirtzoglou, E., Panayiotidis, M. I., Galanis, A. and Plessas, S. (2018).** Production of a novel functional fruit beverage consisting of cornelian cherry juice and probiotic bacteria. *Antioxidants* **7**(11), 163.
- Mohanty, D., Misra, S., Mohapatra, S. and Sahu, P. S. (2018).** Probiotics and synbiotics: Recent concepts in nutrition. *Food Bioscience* **26**, 152-160.
- Mokhtari, S., Jafari, S. M. and Khomeiri, M. (2019).** Survival of encapsulated probiotics in pasteurized grape juice and evaluation of their properties during storage. *Food Science and Technology International* **25**(2), 120-129.
- Moon, K. M., Kwon, E. B., Lee, B. and Kim, C. Y. (2020).** Recent trends in controlling the enzymatic browning of fruit and vegetable products. *Molecules* **25**(12), 2754.
- Mostafa, H. S., Ali, M. R. and Mohamed, R. M. (2021).** Production of a novel probiotic date juice with anti-proliferative activity against Hep-2 cancer cells. *Food Science and Technology* **41**(Suppl 1), 105-115.
- Mubaiwa, J., Fogliano, V., Chidewe, C. and Linnemann, A. R. (2018).** Bambara groundnut (*Vigna subterranea* (L.) Verdc.) flour: A functional ingredient to favour the use of an unexploited sustainable protein source. *PLoS ONE* **13**(10), e0205776.
- Nagpal, M., Maheshwari, D., Rakha, P., Dureja, H., Goyal, S. and Dhingra, G. (2012).** Formulation development and evaluation of alginate microspheres of ibuprofen. *Journal of Young Pharmacists* **4**(1), 13-16.
- Nematollahi, A., Sohrabvandi, S., Mortazavian, A. M. and Jazaeri, S. (2016).** Viability of probiotic bacteria and some chemical and sensory characteristics in cornelian cherry juice during cold storage. *Electronic Journal of Biotechnology* **21**, 49-53.
- Ng, S. L., Lai, K. W., Nyam, K. L. and Pui, L. P. (2019).** Microencapsulation of *Lactobacillus plantarum* 299v incorporated with oligofructose in chitosan coated-alginate beads and its storage stability in amarella juice. *Malaysian Journal of Microbiology* **15**(5), 408-418.
- Nualkaekul, S., Salmeron, I. and Charalampopoulos, D. (2011).** Investigation of the factors influencing the survival of *Bifidobacterium longum* in model acidic solutions and fruit juices. *Food Chemistry* **129**(3), 1037-1044.

- Obulesu, M. and Bhattacharya, S. (2011).** Color changes of tamarind (*Tamarindus indica* L.) pulp during fruit development, ripening, and storage. *International Journal of Food Properties* **14(3)**, 538-549.
- Odriozola-Serrano, I., Soliva-Fortuny, R. and Martín-Belloso, O. (2008).** Phenolic acids, flavonoids, vitamin C and antioxidant capacity of strawberry juices processed by high-intensity pulsed electric fields or heat treatments. *European Food Research and Technology* **228(2)**, 239-248.
- Oladipo, I. C., Atolagbe, O. O. and Adetiba, T. M. (2014).** Nutritional evaluation and microbiological analysis of yoghurt produced from full cream milk, tiger-nut milk, skimmed milk and fresh cow milk. *Pensee Journal* **76(4)**, 30-38.
- Padma Ishwarya, S. and Prabhasankar, P. (2014).** Prebiotics: Application in bakery and pasta products. *Critical Reviews in Food Science and Nutrition* **54(4)**, 511-522.
- Pereira, A. L. F., Maciel, T. C. and Rodrigues, S. (2011).** Probiotic beverage from cashew apple juice fermented with *Lactobacillus casei*. *Food Research International* **44(5)**, 1276-1283.
- Petruzzi, L., Campaniello, D., Speranza, B., Corbo, M. R., Sinigaglia, M. and Bevilacqua, A. (2017).** Thermal treatments for fruit and vegetable juices and beverages: A literature overview. *Comprehensive Reviews in Food Science and Food Safety* **16(4)**, 668-691.
- Phuapaboon, P., Leenanon, B. and Levin, R. E. (2013).** Effect of *Lactococcus lactis* immobilized within pineapple and yam bean segments, and Jerusalem artichoke powder on its viability and quality of yogurt. *Food and Bioprocess Technology* **6(10)**, 2751-2762.
- Rai, A., Das, S., Chamallamudi, M. R., Nandakumar, K., Shetty, R., Gill, M., Sumalatha, S., Devkar, R., Gourishetti, K. and Kumar, N. (2018).** Evaluation of the aphrodisiac potential of a chemically characterized aqueous extract of *Tamarindus indica* pulp. *Journal of Ethnopharmacology* **210**, 118-124.
- Renuka, B., Kulkarni, S. G., Vijayanand, P. and Prapulla, S. G. (2009).** Fructooligosaccharide fortification of selected fruit juice beverages: Effect on the quality characteristics. *LWT - Food Science and Technology* **42(5)**, 1031-1033.
- Rieder, A. and Samuelsen, A. B. (2012).** Do cereal mixed-linked β -glucans possess immune-modulating activities? *Molecular Nutrition and Food Research* **56(4)**, 536-547.
- Sahadeva, R. P. K., Leong, S. F., Chua, K. H., Tan, C. H., Chan, H. Y., Tong, E. V., Wong, S. Y. W. and Chan, H. K. (2011).** Survival of commercial probiotic strains to pH and bile. *International Food Research Journal* **18(4)**, 1515-1522.
- Santhirasegaram, V., Razali, Z. and Somasundram, C. (2013).** Effects of thermal treatment and sonication on quality attributes of Chokanan mango (*Mangifera indica* L.) juice. *Ultrasonics Sonochemistry* **20(5)**, 1276-1282.
- Savedboworn, W., Niyomrat, S., Nakovn, J. and Phattayakorn, K. (2017).** Impact of inulin on viability and storage stability of probiotic *Lactobacillus plantarum* TISTR 2075 in fermented rice extract. *Agriculture and Natural Resources* **51(6)**, 463-469.
- Savedboworn, W., Teawsomboonkit, K., Surichay, S., Riansa-ngawong, W., Rittisak, S., Charoen, R. and Phattayakorn, K. (2019).** Impact of protectants on the storage stability of freeze-dried probiotic *Lactobacillus plantarum*. *Food Science and Biotechnology* **28(3)**, 795-805.
- Sheehan, V. M., Ross, P. and Fitzgerald, G. F. (2007).** Assessing the acid tolerance and the technological robustness of probiotic cultures for fortification in fruit juices. *Innovative Food Science and Emerging Technologies* **8(2)**, 279-284.
- Shi, L. H., Balakrishnan, K., Thiagarajah, K., Mohd Ismail, N. I. and Yin, O. S. (2016).** Beneficial properties of probiotics. *Tropical Life Sciences Research* **27(2)**, 73-90.
- Siang, S. C., Wai, L. K., Lin, N. K. and Phing, P. L. (2019).** Effect of added prebiotic (isomalto-oligosaccharide) and coating of beads on the survival of microencapsulated *Lactobacillus rhamnosus* GG. *Food Science and Technology* **39(Suppl 2)**, 601-609.
- Stack, H. M., Kearney, N., Stanton, C., Fitzgerald, G. F. and Ross, R. P. (2010).** Association of beta-glucan endogenous production with increased stress tolerance of intestinal lactobacilli. *Applied and Environmental Microbiology* **76(2)**, 500-507.
- Suri, S., Kumar, V., Prasad, R., Tanwar, B., Goyal, A., Kaur, S., Gat, Y., Kumar, A., Kaur, J. and Singh, D. (2019).** Considerations for development of lactose-free food. *Journal of Nutrition and Intermediary Metabolism* **15**, 27-34.
- Tril, U., Fernández-López, J., Álvarez, J. Á. P. and Viuda-Martos, M. (2014).** Chemical, physicochemical, technological, antibacterial and antioxidant properties of rich-fibre powder extract obtained from tamarind (*Tamarindus indica* L.). *Industrial Crops and Products* **55**, 155-162.
- Yee, W. L., Yee, C. L., Lin, N. K. and Phing, P. L. (2019).** Microencapsulation of *Lactobacillus acidophilus* NCFM incorporated with mannitol and its storage stability in mulberry tea. *Ciência e Agrotecnologia* **43**, e005819.
- Yeung, T. W., Üçok, E. F., Tiani, K. A., McClements, D. J. and Sela, D. A. (2016).** Microencapsulation in alginate and chitosan microgels to enhance viability of *Bifidobacterium longum* for oral delivery. *Frontiers in Microbiology* **7**, 494.