

“EFFECT OF PHILHEALTH ENROLMENT IN THE DELAY IN DISCHARGE OF PEDIATRIC PATIENTS WITH ACUTE ILLNESS IN A GOVERNMENT TERTIARY HOSPITAL”

JAN KAMILLE R. CORONEL, MD AND SORAYA ALVARADO, MD

ABSTRACT

Background: Philhealth, the national health insurance program, aims to reduce out-of-pocket expenditure to eradicate blockage to healthcare access by subsidizing expenses during admission. A possible determinant of its success is the timeliness of management and prevention of discharge delay.

Objective: To determine the association of Philhealth enrollment with the timeliness of patient discharge in a government tertiary hospital.

Methods: A retrospective cohort study involving 98 patients (49 Philhealth members and 49 non-Philhealth members) was done. Data including age, sex, admitting diagnosis, social service classification, and type of Philhealth membership were collated from the admitting section. Reasons for delay in discharge and number of days delayed was obtained from discharge clearance list of the social service unit. Association between PhilHealth coverage and hospital delay was analyzed using logistic regression analysis.

Results: Discharge delay is 3 times more likely to occur among non-Philhealth members as compared to the Philhealth group. Other variables were not significantly associated with discharge delay.

Conclusions: Discharge delay is significantly associated with non-membership to Philhealth. Further investigation through focus group discussions with patients’ families may be done to uncover other possible causes for discharge delay.

INTRODUCTION

Hospital length of stay has been used as an indicator of efficiency of hospital resource utilization [4, 14]. Although it is largely affected by disease severity and several other non-patient factors, it reflects how proficiently hospital processes are carried out from the time of admission to discharge. Any delay in such processes ultimately leads to prolonged hospital stay.

Financial barrier is only one of the factors identified to contribute to extension of patients’ hospital stay. In the Philippine setting, it may be deemed as the major consideration for delays in discharge. The “fear of incurring a significant financial burden for health care or falling into a ‘medical poverty trap’ can cause ill individuals to either delay or refrain from seeking essential medical services “ [11]. The same rationale may explain to some extent, the

delays encountered in performing necessary procedures and administering essential medications in the in-patient setting. Patients who delay consult for any condition are more likely to present with more severe manifestations and complications as compared to those who seek healthcare early into the disease.

The National Health Insurance Program embodied by the Philippine Health Insurance Corporation (Philhealth), is the government’s effort to have a universal healthcare coverage for all Filipinos in order to eliminate finances as a stumbling block for health utilization. Since Philhealth can cover the expenses for laboratory work-up and medications, utilization of its benefits reduces the financial burden of patients. In so doing, it also speeds up the diagnostic and therapeutic management of admitted patients leading to faster recovery and timely discharge. However, since majority of the health

expenditure still comes from out-of-pocket resources, the effect of having Philhealth coverage on the health outcomes is yet to be established. Therefore, this study aims to determine the association of Philhealth benefit availment with the timeliness of discharge of patients admitted for acute conditions.

The results of this study can help determine whether enrolment of patients to Philhealth contribute to maximization of the hospitals resources, particularly bed turnover. Also, this study may be able to elucidate other bottlenecks in the discharge planning process,

We aimed to determine the effect of Philhealth enrollment in the timeliness of discharge of patients with acute illness in our institution. Our specific objectives were:

- 1) To present the demographic profile of patients who have utilized Philhealth benefits during their admission
- 2) To demonstrate how many patients stayed more than 1 day after being given a send home order
- 3) To determine how many patients who have Philhealth accounts overstayed
- 4) To identify other reasons for delay in discharge of patients with acute illness
- 5) To illustrate the current prevalence of Philhealth coverage in the Philippines and compare it to that in our institution

METHODOLOGY

A retrospective cohort study was done to determine which among those children admitted for acute illness were enrolled and availed of Philhealth benefits and were discharged timely. The sample population was randomly selected from the database of admitted patients from January 2016 to June 2016. All private patients were excluded. Those patients who have chronic illness as co-morbidities (i.e. all forms of malignancy, cardiac conditions, neurologic conditions) as well as neonates were also excluded. However, those patients admitted for acute exacerbations of a chronic illness are included (i.e. bronchial asthma).

Using NCSS-PASS 2013, the minimum sample size requirement is at least 98 based on the percentage of uninsured patients with hospital delay = 32.2% [15] and odds ratio= 4.51 [12] with level of significance= 5% and power= 80%.

The service patients were classified as to having Philhealth membership or not. From the list of patients with Philhealth, 49 were randomly selected. The same number was randomly selected from the list of non-Philhealth patients. All patients who are enrolled at the beginning of the study, except those who expired, were analyzed at the end of the study regardless of outcome (i.e. acute illness which complicated during admission).

The demographic profile (age, sex), the acute condition for which the patient was admitted, the social service classification and the Philhealth membership type were taken from the patient database from the Admitting section. The discharge delay form was duly accomplished based on such data.

A list of discharge clearances during the time of the study was obtained from the Social Services Unit. From such list, each patient included in the study was cross-referenced to determine delay in discharge. If the patient was not in the list, the patient was considered able to go home on the day the discharge order was made. On the other hand, if the patient's name is on the discharge clearance list, the number of discharge clearances was noted. This was noted in the discharge delay form. The number of discharge clearances was equivalent to the number of days that the patient's discharge was delayed from the time the order for discharge was made. The reason for the delay is also noted in the discharge clearance list provided by the Social Service Unit. Moreover, fund sources (PCSO, DSWD, etc.) for settling the hospital bill of patients who were unable to go home at the discharge point were recorded in the said document.

The collected data were tabulated in Microsoft Excel. The characteristics of the sample population like age, sex, social service


classification (C1, C2 and C3), diagnosis and type of Philhealth membership (for the Philhealth group) were analyzed against the timeliness of discharge. The association between variables was determined using the chi-square test.

Data analysis was performed in Stata SE version 13. Quantitative variables were summarized as mean and standard deviation, while qualitative variables were tabulated as frequency and percentage. Comparison of baseline characteristics was analyzed using independent t-test for quantitative variables, and Fisher’s exact test for qualitative variables. Association between PhilHealth coverage and hospital delay was analyzed using logistic regression analysis. The level of significance was set at 5%.

RESULTS

There are 3,302 patients who were admitted at the service wards from January – June 2016. Of these patients, 2,845 or 86% were enrolled to Philhealth, while the remaining 457 or 14% were non-Philhealth members. There was a total of 98 patients included in the study, 49 are enrolled in Philhealth while the remaining are non-Philhealth members. Approximately 45% of the patients belong to the 1 – 4 years old age group. Majority of the patients included in the study are females, comprising 62%. The most common conditions of patients included in the study are pneumonia (32%), acute gastroenteritis (11%) and acute glomerulonephritis (9%) (Figure 1). This finding is reflective of the hospital-wide data, which also shows these conditions to be the most prevalent diagnoses of admitted patients.

Figure 1. Acute Conditions of Patients in the Study.



Based on the social service classification, most of the patients (75%) are classified as C3, indicating that a considerable number of patients belong to the lower income bracket (Table 1).


Table 1. Demographic Profile of Subjects with Philhealth and Non-Philhealth

	With Philhealth (n = 49)	Without Philhealth (n = 49)	p-value
Age			
1 mo – 11 mos	13 (26.53%)	15 (30.61%)	0.332
1 – 4 years old	26 (53.06%)	18 (36.73%)	
5 – 9 years old	4 (8.16%)	10 (20.41%)	
10 – 13 years old	3 (6.12%)	4 (8.16%)	
14 – 18 years old	3 (6.12%)	2 (4.08%)	
Sex			
Male	19 (38.78%)	18 (36.73%)	1.00
Female	30 (61.22%)	31 (63.27%)	
Social Service Classification			
C1	0 (0%)	1 (2.04%)	0.152
C2	15 (30.61%)	8 (16.33%)	
C3	34 (69.39%)	40 (81.63%)	

However, it is of note that there are more patients in the C3 group who do not have Philhealth. This contrasts with the other social service classification group (C2), which shows that there are more patients who are enrolled to Philhealth than those who are non-members. Further classification of the Philhealth group

based on the membership type showed that the most common membership type is the SSS-dependent type comprising 51% of the said group (Figure 2), while indigent-dependent members and those who have voluntary contributions consist 40% and 9%, respectively.

Figure 2. Distribution of Patients with Philhealth based on Membership Type



The characteristics of enrolled patients such as age, sex and acute illness for which they are admitted were analyzed against discharge delay to identify other contributory factors, but none of them demonstrated significant association. However, analyzing Philhealth enrolment against delay in discharge showed that non-membership to Philhealth is a risk

factor for delay in discharge, in that those without Philhealth are 3 - 4 times more likely to experience delay in discharge compared to those with Philhealth (Table 2). On the average, those who experience delay in discharge stay in the hospital for 1-2 days after being given the discharge order.

Table 2. Discharge Delay among Study Patients.

	With delay (n = 16)	Without Delay (n = 82)	P-value
Age			
1 mo – 11 mos	3 (18.75%)	25 (30.5%)	0.08
1 – 4 years old	9 (56.3%)	35 (42.7%)	
5 – 9 years old	0 (0%)	14 (17.1%)	
10 – 13 years old	3 (18.75%)	4 (4.88%)	
14 – 18 years old	1 (6.25%)	4 (4.88%)	
Sex			
Male	7 (43.7%)	30 (36.6%)	0.58
Female	9 (56.3%)	52 (63.4%)	
Acute Illness			
Abscess	0 (0%)	1 (1.22%)	0.59
Acute Gastritis	0 (0%)	3 (3.66%)	
Acute Glomerulonephritis	0 (0%)	9 (10.98%)	
Acute Gastroenteritis	2 (12.5%)	9 (10.98%)	
Kawasaki Disease	0 (0%)	5 (6.10%)	
Bronchial Asthma	0 (0%)	4 (4.88%)	
Febrile Convulsions	3 (18.75%)	4 (4.88%)	
Bronchiolitis	0 (0%)	2 (2.44%)	
Candidiasis	0 (0%)	1 (1.22%)	
Cellulitis	0 (0%)	1 (1.22%)	
Central Nervous System (CNS) Infection	0 (0%)	3 (3.66%)	
Sepsis	1 (6.25%)	1 (1.22%)	
Dengue	1 (6.25%)	4 (4.88%)	
Type I Diabetes Mellitus	1 (6.25%)	2 (2.44%)	
Fever of Unknown Origin	0 (0%)	1 (1.22%)	
Hypokalemic Periodic Paralysis	1 (6.25%)	0 (0%)	
Henoch-Schonlein Purpura			
Idiopathic Thrombocytopenic Purpura	0 (0%)	1 (1.22%)	
Pneumonia	0 (0%)	3 (3.66%)	
Scabies			
Traumatic Brain Injury	7 (43.75%)	25 (30.49%)	
	0 (0%)	1 (1.22%)	
	0 (0%)	2 (2.44%)	
Philhealth Membership			
With Philhealth	4 (25%)	45 (54.9%)	0.03
Without Philhealth	12 (75%)	37 (45.1%)	

All patients who experienced delay in discharge identified financial constraint as the reason for delay. The patients' discharges were facilitated by securing funds out-of-pocket and from government agencies such as the Philippine Charity Sweepstakes Office (PCSO) and Department of Social Welfare and Development (DSWD). For the Philhealth

group, all patients who experienced delay utilized out-of-pocket resources to fund their hospital bill. Similar findings were seen in the non-Philhealth group - 83% of those who had delay in discharge used out-of-pocket resources to pay for their hospital bill. Other sources of funding identified in this study were PCSO and DSWD (Table 3).

Table 3. Non-Philhealth Sources of Funding Among Patients with Delay

Source of Funding	With Philhealth (n = 4)	Without Philhealth (n = 12)
Out-of-pocket	4 (100%)	10 (83%)
PCSO	0 (0%)	1 (8.3%)
DSWD	0 (0%)	1 (8.3%)

DISCUSSION

Delay in discharge contributes to the cascade of administrative concerns of hospitals, especially in the setting of resource scarcity. Discharge delays have been shown to increase the cost of treatment and even aggravate the patients’ outcome (15). In the study by Gaughan et al., the cost of delay in discharge is identified as both financial and clinical. The hospital tends to unnecessarily spend resources for a patient who is deemed clinically fit for discharge by allotting a bed which could have been occupied by another patient who has a greater need for medical service. Its clinical implication is also note-worthy as patients who experience delay in discharge may acquire hospital-associated infections, leading to a prolonged hospital stay than expected.

Studies have been made to elucidate the factors contributing to discharge delays to adequately address them. Most researchers conclude that majority of discharges are delayed due to medical causes (awaiting laboratory results, managing complications, awaiting consultants’ opinion). However, little is known regarding non-medical causes of discharge delay, moreso in the Philippine setting.

Financial capacity to access healthcare has been identified in several studies to cause delay in seeking treatment (11, 16). Social health insurance health programs have been developed and encouraged to alleviate the financial burden that has been an impediment to healthcare access, especially by the poor. In the study by Fowler et al., being uninsured was identified as a risk factor for discharge delay presumably because these patients arrive in the hospital in a more critical state, needing more intensive management due to delay in initial medical

consult. However, the direct effect of health insurance status and its utilization to health outcomes has not been thoroughly considered, especially in the local setting.

An indicator of a successful national health insurance program is a decreased percentage of out-of-pocket expenditure (16). Therefore, reinforcing the national health insurance program through Philhealth benefit utilization is one strategy to alleviate the financial concerns of the patients and expedite the diagnosis, management and ultimately the discharge of these patients.

During the study period, 86% of admitted patients are Philhealth members while the remaining are non-Philhealth members. This is comparable to the statistics released by Philhealth where approximately 9 out of 10 Filipinos are covered by Philhealth. By increasing their coverage, Philhealth comes close to achieving its aim to for universal healthcare by easing the financial burden imposed by healthcare costs. Aside from voluntary and compulsory contributions, the government also subsidizes funding of the said program especially for those belonging to the lowest economic quintiles through the Indigent Program (IP). Despite these efforts, there are still many Filipinos who remain non-members or become members but do not utilize the benefits of Philhealth to avail healthcare services. In the 2015 Philhealth statistics, only 12% of total eligible Philhealth members utilize its benefits. In this research, however, it is assumed that Philhealth membership is equivalent to utilization of its benefits during the admission. Philhealth members who did not avail of Philhealth benefits were not identified in this paper.

The reasons for non-enrolment to Philhealth are beyond the scope of this study. In a study by Faraon et al., sex, income and type of Philhealth membership were identified variables to be significantly associated with underutilization of Philhealth. The outcome of this study had a similar result. Based on the patients' social service classification, vis-à-vis their income status, most of non-Philhealth members belong to C3 or the lowest income group. Such finding may be supposed to be also linked to the educational attainment and subsequently, the knowledge and awareness of the families of these patients regarding the Philhealth benefits and how to avail them. Other reasons for underutilization of Philhealth are as follows:

- Lack of knowledge on benefits
- Lack of knowledge on filing claims
- Lack of cooperation from doctors in getting required documents
- Cumbersome or unmanageable process
- High transaction costs compared to benefits
- Ineligible claim

Although the association between the social service classification and Philhealth membership was not found to be significant in this study, it may be prudent to consider such pattern in succeeding investigations.

In this study, most patients are subscribed to Philhealth by compulsory contributions through the employer, also known as SSS-dependent members, comprising 51% of the sample population. However, Latest Philhealth data demonstrates that majority of covered members are indigents, comprising 49%. This discrepancy may be explained by the fact that some patients are initially admitted as non-Philhealth members and are only enrolled to Philhealth during admission, through the assistance of the social service unit. These patients are usually indigents who are granted a no balance billing status (NBB) during their stay. Such enrolment mechanism is noted in the records of social service and the admitting section. However, none of the patients who are included in this study experienced said

mechanism of enrolment to Philhealth. All non-Philhealth members remained unenrolled to Philhealth until discharge.

Those patients who experienced delay in discharge reported financial setback as the reason for the delay. Data from the social service unit showed that majority of these patients still resorted to out-of-pocket resources to settle their hospital bill, regardless of their Philhealth membership. However, there are 2 patients who sought financial assistance from PCSO and DSWD. This agrees with the finding that household budget for healthcare is still mostly out-of-pocket (16). This reliance on out-of-pocket sources for healthcare has detrimental effects. In the study by Ulep and Dela Cruz, high out-of-pocket expenditure for health needs can:

- redistribute income “in the wrong direction” (i.e., from chronically ill to healthy individuals and, typically, from the relatively poor to the relatively affluent groups [Plumper and Neumayer 2012])
- lead people to make tough choices concerning their health such as not complying with prescribed drug use due to high costs, forgoing necessities, or borrowing money to pay for prescriptions;
- affect women and minorities who may forgo critical prevention screenings and skimp on medications due to high costs; and
- increase the financial burden on those with valid insurance (Aji et al. 2013)

CONCLUSION

In summary, this study demonstrated that Philhealth enrolment and its subsequent utilization is associated with timeliness of discharge of pediatric patients admitted for acute illnesses. Non-membership to the national health insurance program increases the risk of discharge delay three-fold. Other patient factors such as age, sex, social service classification and the acute illness for which the patients are admitted did not show any significant association with discharge delay.

Further investigations can be made to elucidate on other factors affecting discharge delay. This study is limited to review of patients' records from the admitting section and social service unit. It is recommended that a prospective study be done to identify the reasons for delay in discharge as will be reported by patients' families, which may not be solely financial in nature. Focus group discussions among the patients' parents may be done to fully understand the reasons for delay. Such method of study can also give light to the reasons for non-Philhealth membership. Recognizing other variable affecting discharge delay may be helpful for the hospital administration and even the national government, to draft policies and systems to improve the health insurance program so as to achieve universal healthcare for all.

REFERENCES:

- 1) Mawajdeh, S., et al. The Effect of Type of Hospital and Health Insurance on Hospital Length of Stay in Irbid, North Jordan. *Health and Policy Planning (1997) 12(2), 166-172.*
- 2) Sung-tse, Li, et al. Factors Affecting Length of Stay in the Pediatric Emergency Department. *Pediatrics and Neonatology (2013)54, 179- 187*
- 3) Romano, P. et al. Selecting Quality and Resource Use Measures: A Decision Guide for Community Quality Collaboratives. *Agency for Healthcare Research and Quality. Publication No. 09(10)-0073 May 2010*
- 4) "Average length of stay in hospitals", in Health at a Glance 2011: OECD Indicators, OECD Publishing. <http://dx.doi.org/10.1787/health_glance-2011-33-en> Date Accessed: June 27, 2015.
- 5) Hendy, P. et al. "In-depth analysis of delays to patient discharge: a metropolitan teaching hospital experience". *Clinical Medicine 2012, Vol 12, No 4: 320-3.*
- 6) Tak, S. et al. "A comparative time motion study of all types of patient discharges in a hospital". *Global Journal of Medicine and Public Health.*
- 7) Quimbo, S. et al. "Underutilization of Social Insurance among the Poor: Evidence from the Philippines". 2008. *PLoS ONE 3(10): e3379.doi:10.1371/journal.pone.0003379*
- 8) Puyat, M.E. "Factors Influencing PhilHealth Coverage and In-patient Benefit Utilization of Filipino Children under Five". Philippine Institute for Development Studies. Discussion Paper Series no. 2013-37.
- 9) Kraft, A. et al. "The Health and Cost Impact of Care Delay and the Experimental Impact of Insurance on Delays: Evidence from a Developing Country". *Journal of Pediatrics. 2009 August ; 155(2): 281-5.e1. doi:10.1016/j.jpeds.2009.02.035.*
- 10) Tobe, M. et al. "Out-of-pocket medical expenses for inpatient care among beneficiaries of the National Health Insurance Program in the Philippines". *Health Policy and Planning 2012;1-13. doi:10.1093/heapol/czs092*
- 11) Fowler, R. et al. "An Official American Thoracic Society Systematic Review: The Association between Health Insurance Status and Access, Care Delivery, and Outcomes for Patients Who Are Critically Ill". *American Thoracic Society Documents. October 2009.*
- 12) Freitas, A. "Factors Influencing Hospital High Length of Stay". *BMC Health Services Research 2012, 12:265 <<http://www.biomedcentral.com/1472-6963/12/265>>. Date Accessed: November 8, 2015.*

- 13) Kulinskaya, E., et al. Length of stay as a performance indicator: robust statistical methodology. *IMA Journal of Management Mathematics* (2005) 16, 369–381.
- 14) Cooper, K. “Delayed Hospital Discharge Figures still Rising”. British Medical Association. <www.bma.org.uk>. Date Accessed: July 7, 2016
- 15) Gaughan, J. et al. “Delayed Discharges and Hospital Type: Evidence from the English NHS”. CHE Research Paper 133. University of York, July 2016.
- 16) Ulep, V. and Dela Cruz, N. “Analysis of Out-of-Pocket Expenditures in the Philippines”. *Philippine Journal of Development*. Number 72, First and Second Semesters 2013 Volume XL, Numbers 1 &2
- 17) Faraon, E. et al. “Significant Predictors of Underutilization of Inpatient Benefits among PhilHealth Members in Selected Barangays in Manila”. *Acta Medica Philippina*. Vol. 47 No. 3, 2013.
- 18) Senate Economic Planning Office (SEPO). “Philhealth at a Glance”. November 2009.
- 19) Philhealth: 2015 Stats and Charts. <philhealth.gov.ph>. Date Accessed: October 1, 2016