



OFFICIAL PUBLICATION OF
THE MINISTRY OF HEALTH
AND
UNIVERSITI BRUNEI DARUSSALAM

Brunei International Medical Journal

Volume 19

3 November 2023 (19 Rabiulakhir 1445H)

EVALUATION OF QUALITY OF LIFE IN PATIENTS WITH A FACIAL PROSTHESIS.

Song Shan HUI, Normastura AB RAHMAN, Rabihah ALAWI, Nor Aidaniza ABDUL MUTTLIB.

School of Dental Sciences, Universiti Sains Malaysia, Kampus Kesihatan, 16150
Kubang Kerian, Kelantan, Malaysia.

ABSTRACT

Introduction: This study aimed to assess the quality of life of patients with facial prostheses after head and neck surgery. **Materials and Methods:** Sixty-four patients who had received an extraoral or facial prosthesis over the previous five years and had been using it for at least six months after the treatment were interviewed. A validated questionnaire, the University of Washington Quality of Life Questionnaire version 4, (UoW- QOL v4), was used and the patients' sociodemographic profiles were obtained. Data was analysed using SPSS ver. 22. **Results:** A total of 64 patients, with a mean age of 38.1 years, were included, 41 (64.1%) of whom were male and 23 (35.9%) were female. The most significant problems encountered by the patients were appearance (14.1%), mood (11%) and recreation (7.8%). The majority stated that their overall quality of life was 'fair' or 'good' (84.4%). Pain (90.6%), activity (87.5%), and appearance (78.1%) were the issues most frequently reported by patients. **Conclusion:** The quality-of-life scores, as assessed using the (UoW-QOL v4) scale, were fair or good in patients with facial prostheses. Pain, activity, and appearance were regarded as the most important issues affecting their quality of life.

Keywords: Evaluation, Head and neck cancer, Maxillofacial prosthesis, Quality of life, UoW-QOL v4.

Brunei Int Med J. 2023;19:63-69

Brunei International Medical Journal (BIMJ)

Official Publication of The Ministry of Health and Universiti Brunei Darussalam

EDITORIAL BOARD

Editor-in-Chief	Ketan PANDE
Sub-Editors	Vui Heng CHONG William Chee Fui CHONG
Editorial Board Members	Muhd Syafiq ABDULLAH Alice Moi Ling YONG Ahmad Yazid ABDUL WAHAB Jackson Chee Seng TAN Pemasiri Upali TELISINGHE Pengiran Khairol Asmee PENGIRAN SABTU Dayangku Siti Nur Ashikin PENGIRAN TENGAH

INTERNATIONAL EDITORIAL BOARD MEMBERS

Lawrence HO Khek Yu (Singapore)	Chuen Neng LEE (Singapore)
Wilfred PEH (Singapore)	Emily Felicia Jan Ee SHEN (Singapore)
Surinderpal S BIRRING (United Kingdom)	Leslie GOH (United Kingdom)
John YAP (United Kingdom)	Ian BICKLE (United Kingdom)
Nazar LUQMAN (Australia)	Christopher HAYWARD (Australia)
Jose F LAPENA (Philippines)	

Advisor

Wilfred PEH (Singapore)

Past Editors-in-Chief

Nagamuttu RAVINDRANATHAN
Kenneth Yuh Yen KOK
Chong Vui Heng
William Chong Chee Fui

Proof reader

John WOLSTENHOLME (CfBT Brunei Darussalam)

Aim and Scope of Brunei International Medical Journal

The Brunei International Medical Journal (BIMJ) is a six-monthly peer-reviewed official publication of the Ministry of Health under the auspices of the Clinical Research Unit, Ministry of Health, Brunei Darussalam.

The BIMJ publishes articles ranging from original research papers, review articles, medical practice papers, special reports, audits, case reports, images of interest, education and technical/innovation papers, editorials, commentaries, and letters to the Editor. Topics of interest include all subjects that relate to clinical practice and research in all branches of medicine, basic and clinical including topics related to allied health care fields. The BIMJ welcomes manuscripts from contributors but usually solicits review articles and special reports. Proposals for review papers can be sent to the Managing Editor directly. Please refer to the contact information of the Editorial Office.

INSTRUCTION TO AUTHORS

Manuscript submissions

All manuscripts should be sent to the Managing Editor, BIMJ, Ministry of Health, Brunei Darussalam; e-mail: bimjonline@gmail.com. Subsequent correspondence between the BIMJ and authors will, as far as possible be conducted via email quoting the reference number.

Conditions

Submission of an article for consideration for publication implies the transfer of the copyright from the authors to the BIMJ upon acceptance. The final decision of acceptance rests with the Editor-in-Chief. All accepted papers become the permanent property of the BIMJ and may not be published elsewhere without written permission from the BIMJ.

Ethics

Ethical considerations will be taken into account in the assessment of papers that have experimental investigations of human or animal subjects. Authors should state clearly in the Materials and Methods section of the manuscript that the institutional review board has approved the project. Those investigators without such review boards should ensure that the principles outlined in the Declaration of Helsinki have been followed.

MANUSCRIPT CATEGORIES

Original articles

These include controlled trials, interventional studies, studies of screening and diagnostic tests, outcome studies, cost-effectiveness analyses, and large-scale epidemiological studies. The manuscript should include the following; introduction, materials

and methods, results, and conclusion. The objective should be stated clearly in the introduction. The text should not exceed 2500 words and references not more than 30.

Review articles

These are, in general, invited papers, but unsolicited reviews, if of good quality, may be considered. Reviews are systematic critical assessments of literature and data sources on clinical topics, emphasising factors such as cause, diagnosis, prognosis,

therapy, or prevention. Reviews should be made relevant to our local setting and preferably supported by local data. The text should not exceed 3000 words and references not more than 40.

Special Reports

This section usually consists of invited reports that have a significant impact on healthcare practice and usually cover disease outbreaks, management guidelines, or policy statement papers.

Audits

Audits of relevant topics generally follow the same format as the original article and the text should not exceed 1,500 words and references not more than 20.

Case reports

Case reports should highlight interesting rare cases or provide good learning points. The text should not exceed 1000 words; the number of tables, figures, or both should not be more than two, and references should not be more than 15.

Education section

This section includes papers (i.e. how to interpret ECG or chest radiography) with the particular aim of broadening knowledge or serving as revision materials. Papers will usually be invited but well-written papers on relevant topics may be accepted. The text should not exceed 1500 words and should include not more than 15 figures illustrations and references should not be more than 15.

Images of interest

These are papers presenting unique clinical encounters that are illustrated by photographs, radiographs, or other figures. The image of interest should include a brief description of the case and a discussion of educational aspects. Alternatively, a mini quiz can be presented and answers will be posted in a different section of the publication. A maximum of three relevant references should be included. Only images of high quality (at least 300 dpi) will be acceptable.

Technical innovations

This section includes papers looking at novel or new techniques that have been developed or introduced to the local setting. The text should not exceed 1000 words and should include not more than 10 figures illustrations and references should not be more than 10.

Letters to the Editor

Letters discussing a recent article published in the BIMJ are welcome and should be sent to the Editorial Office by e-mail. The text should not exceed 250 words; have no more than one figure or table, and five references.

Criteria for manuscripts

Manuscripts submitted to the BIMJ should meet the following criteria: the content is original; the writing is clear; the study methods are appropriate; the data are valid; the conclusions are reasonable and supported by the data; the information is important; and the topic has a general medical interest. Manuscripts will be accepted only if both their contents and style meet the standards required by the BIMJ.

Authorship information

Designate one corresponding author and provide a complete address, telephone and fax numbers, and e-mail address. The number of authors of each paper should not be more than twelve; a greater number requires justification. Authors may add a publishable footnote explaining the order of authorship.

Group authorship

If authorship is attributed to a group (either solely or in addition to one or more individual authors), all members of the group must meet the full criteria and requirements for authorship described in the following paragraphs. One or more authors may take responsibility 'for' a group, in which case the other group members are not authors, but may be listed in an acknowledgment.

Authorship requirement

When the BIMJ accepts a paper for publication, authors will be asked to sign statements on (1) financial disclosure, (2) conflict of interest, and (3) copyright transfer. The correspondence author may sign on behalf of co-authors.

Authorship criteria and responsibility

All authors must meet the following criteria: to have participated sufficiently in the work to take public responsibility for the content; to have made substantial contributions to the conception and de-

sign, and the analysis and interpretation of the data (where applicable); to have made substantial contributions to the writing or revision of the manuscript; and to have reviewed the final version of the submitted manuscript and approved it for publication. Authors will be asked to certify that their contribution represents valid work and that neither the manuscript nor one with substantially similar content under their authorship has been published or is being considered for publication elsewhere, except as described in an attachment. If requested, authors shall provide the data on which the manuscript is based for examination by the editors or their assignees.

Financial disclosure or conflict of interest

Any affiliation with or involvement in any organisation or entity with a direct financial interest in the subject matter or materials discussed in the manuscript should be disclosed in an attachment. Any financial or material support should be identified in the manuscript.

Copyright transfer

In consideration of the action of the BIMJ in reviewing and editing a submission, the author/s will transfer, assign, or otherwise convey all copyright ownership to the Clinical Research Unit, RIPAS Hospital, Ministry of Health if such work is published by the BIMJ.

Acknowledgments

Only persons who have made substantial contributions but who do not fulfill the authorship criteria should be acknowledged.

Accepted manuscripts

Authors will be informed of acceptances and accepted manuscripts will be sent for copyediting. During copyediting, there may be some changes made to accommodate the style of the journal format. Attempts will be made to ensure that the overall meaning of the texts is not altered. Authors will be informed by email of the estimated time of publication. Authors may be requested to provide raw data, especially those presented in graphs such as bar charts or figures so that presentations can be constructed following the format and style of the journal. Proofs will be sent to authors to check for any mistakes made during copyediting. Authors are usually given 72 hours to return the proof. No response will be taken as no further corrections are required. Corrections should be kept to a minimum. Otherwise, it may cause a delay in publication.

Offprint

Contributors will not be given any offprint of their published articles. Contributors can obtain an electronic reprint from the journal website.

DISCLAIMER

All articles published, including editorials and letters, represent the opinion of the contributors and do not reflect the official view or policy of the Clinical Research Unit, the Ministry of Health, or the institutions with which the contributors are affiliated to unless this is clearly stated. The appearance of the advertisement does not necessarily constitute an endorsement by the Clinical Research Unit or the Ministry of Health, Brunei Darussalam. Furthermore, the publisher cannot accept responsibility for the correctness or accuracy of the advertisers' text and/or claim or any opinion expressed.

EVALUATION OF QUALITY OF LIFE IN PATIENTS WITH A FACIAL PROSTHESIS.

Song Shan HUI, Normastura AB RAHMAN, Rabihah ALAWI, Nor Aidaniza ABDUL MUTLIB.

School of Dental Sciences, Universiti Sains Malaysia, Kampus Kesihatan, 16150 Kubang Kerian, Kelantan, Malaysia.

ABSTRACT

Introduction: This study aimed to assess the quality of life of patients with facial prostheses after head and neck surgery. **Materials and Methods:** Sixty-four patients who had received an extraoral or facial prosthesis over the previous five years and had been using it for at least six months after the treatment were interviewed. A validated questionnaire, the University of Washington Quality of Life Questionnaire version 4, (UoW- QOL v4), was used and the patients' socio-demographic profiles were obtained. Data was analysed using SPSS ver. 22. **Results:** A total of 64 patients, with a mean age of 38.1 years, were included, 41 (64.1%) of whom were male and 23 (35.9%) were female. The most significant problems encountered by the patients were appearance (14.1%), mood (11%) and recreation (7.8%). The majority stated that their overall quality of life was 'fair' or 'good' (84.4%). Pain (90.6%), activity (87.5%), and appearance (78.1%) were the issues most frequently reported by patients. **Conclusion:** The quality-of-life scores, as assessed using the (UoW-QOL v4) scale, were fair or good in patients with facial prostheses. Pain, activity, and appearance were regarded as the most important issues affecting their quality of life.

Keywords: Evaluation, Head and neck cancer, Maxillofacial prosthesis, Quality of life, UoW- QOL v4.

INTRODUCTION

In overcoming weaknesses with surgical reconstructions, prosthetic rehabilitation requires different approaches. This is the most cost-effective process, allowing short periodic cleansing and the reconstruction of the affected area within a predictable execution period.¹ Due to various specific psychophysical patient

conditions, facial reconstruction using prosthetic rehabilitation is an alternative to surgical reconstruction.²

Patients with facial deformities have been shown to have low self-esteem, which may affect their quality of life.³ Patient functions and self-esteem were reported to improve with the aid of prosthetic rehabilitation.⁴ The effects of the surgery, which include morphological and functional disturbances, were among the criteria to be considered in determining the need for a therapeutic approach in

Corresponding author: Dr. Nor Aidaniza binti Abdul Muttlib, Lecturer in the School of Dental Sciences, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia
T.: +609 7675840, E.: aidaniza@usm.my

patients with post-surgery facial defects.⁵ Surgical reconstruction itself is generally more difficult and does not always produce the desired result.⁶

Problems with the retention of the prosthesis, colour inconsistency, and skin reactions to adhesives and soft tissue layers are among the difficulties reported when facial prostheses are used.⁷ It was reported that within the first three years of service, patient satisfaction with external prostheses decreased.⁸ This was probably due to colour changes and the pigmentation of the prostheses, which compromised their aesthetic qualities.⁹

Ocular prostheses (Figure 1) are made using Polymethylmethacrylate (PMMA), while skin-like prostheses are made from a silicon material. Traditionally, skin adhesives, solvents, eyeglasses, the use of hard and soft tissue undercuts, and other modalities have been used to help in retaining prostheses. Unfortunately, these methods have been associated with weaknesses, for instance, problems with retention, stability, adverse tissue reactions, discolouration and prosthesis deterioration, inconvenience of use or application, poor hygiene, discomfort, and lack of acceptance.¹⁰

Facial prostheses are responsible for



Figure 1: Ocular prostheses.

improving the quality of life of a patient after surgical management of head and neck cancer.⁵ The majority of clinical studies on facial prostheses have reported only biological outcomes. Hence, an evaluation of the effects of prostheses on the quality of life of patients with facial prostheses, using questionnaires developed specifically for this group of patients, may provide important data on treatment outcomes from the patients' perspectives.¹¹ It is important to assess the satisfaction levels and changes in a patient's life that have occurred due to surgery and the rehabilitation process to ensure the prostheses made genuine improvements to overall patient well-being.^{5,12} Hence, this study was conducted to assess the quality of life of patients who had been wearing a facial prosthesis for at least six months but not longer than five years.

METHODOLOGY

A cross-sectional study was conducted among the patients who were using ocular, nasal, or auricular prostheses and who had been rehabilitated at the maxillofacial prosthodontics clinic at Hospital Universiti Sains Malaysia (USM).

The sampling method used in this study was simple random sampling. The reference population was patients who had received facial prostheses at Hospital USM between 2012 and 2018. The sampling was preceded by a review of those patients who had received an extraoral or facial prosthesis over the previous five years and used it for at least six months after the treatment. Sixty-four clinical records were identified. The telephone numbers of patients were recorded and oral consent was taken before they were included in the study.

Patients who were mentally unstable or suffering from dementia, or who had received an intraoral prosthesis, were excluded

from this study.

The data collection involved phone call interviews using a validated questionnaire, the Proforma Questionnaire from the University of Washington Quality of Life Questionnaire version 4, (UoW- QOL v4). The interview sessions took around 20 minutes to complete, on average.

The interview was divided into two parts. The first part referred to the general information about the patients, such as gender, types of prostheses, and the causes of the condition. The second part collected specific data and contained 15 objective questions, including information about the pain, appearance, activity, recreation, speech, chewing, swallowing, shoulder pain, taste, saliva, mood, anxiety, and employment, which were assessed and scored by the patients. The patient had to provide a score from 0-100, whereby 0 was the worst/ poorest score and 100 was the highest/ excellent score. Scores of 80 and above were considered the highest scores, according to Becker *et al.*¹³

Data entry and analysis were done and processed using SPSS software version 24.0. Descriptive statistics were used to summarise and represent the participants' socio-demographic data. Categorical data was represented by frequency and percentage while numerical data was represented by mean and standard deviation.

RESULTS

A total of 100 patients with facial prostheses were available at Hospital USM. Sixty-four patients who fulfilled the inclusion and exclusion criteria were recruited for the study and completed the questionnaire.

Demographic Characteristics

The results showed the respondents have a

mean (SD) age of 38.13 (21.5) and there were more males (n= 41; 64.1%) than females (n=23;35.9%). The majority of reasons why the patients wore facial prostheses were head and neck pathology (n=34; 53.1%), followed by accidents (n=28;43.8%) and congenital conditions (n=2;3.1%).

Types of facial prosthesis

From the data obtained, 61 (95.3%) out of 64 patients were wearing ocular prostheses, 39 (60.9%) of whom were male (Table I). Only 3 (4.7%) patients were wearing auricular prostheses, 2 (3.1%) of whom were male. None of the respondents were wearing nasal or maxillary prostheses.

Assessment of quality of life

Table II represents the summary of three components. Part A concerned the patients' relative feelings a month before their prosthesis's replacement, part B concerned their health-related quality of life during the previous seven days and part C presented the overall quality of life. These were scaled from 0 to 100 to facilitate the presentation, with key results using the same 0 to 100 scale. The findings for part A showed that 51 respondents (79.7%) quoted their health-related quality of life as 'much better' and 13 respondents (20.3%) quoted it as 'somewhat better'. For part B, 50 respondents cited their health-related quality of life during the previous seven days as 'fair' or 'good', while 54 respondents reported their overall quality of life as 'fair' or 'good' for part C.

Table I: Types of facial prostheses (n=64)

Variables	Types of facial prostheses, n (%)			
	Nasal	Auricular	Ocular	Maxillary
Male	0	2 (3.1)	39 (60.9)	0
Female	0	1 (1.6)	22 (34.4)	0
Total	0	3 (4.7)	61 (95.3)	0

Table II: Evaluation of the quality of life

UoW-QOL	Question score									
	0	20	25	40	50	60	75	80	100	
A. Health-related QOL compared to the month before the repalcement	0		0		0		13		51	
B. Health-related QOL during the past 7 days	0	0		23		27		11	3	
C. Overall QOL during the past 7 days	0	0		27		27		7	3	

UoW-QOL: University of Washington Quality of Life

Percentage of patients choosing each domain

Table III shows which domains had been the most important during the previous seven days. Patients were asked to choose up to three domains. Pain (90.6%), activity (87.5%), and appearance (78.1%) were the issues most frequently reported by the patients in this study, followed by recreation (20.3%), anxiety (14.1%), and mood (7.8%). The domains of swallowing, chewing, speech, taste, shoulder pain, and saliva were not chosen by any respondents. This was perhaps because the respondents were wearing extraoral prostheses, which did not interfere with these functions.

Significant problems with each UoW-QOL domain

As shown in Table III, the three domains producing the most significant effects were appearance (14.1%), mood (11%), and recreation (9.7%). The data also suggested that

significant problems existed with other domains but these affected a smaller percentage. Examples of these domains were pain (1.6%), activity (6.3%), and anxiety (3.1%). For the domains of swallowing, chewing, speech, shoulder pain, taste, and saliva, no significant problems were recorded.

DISCUSSION

The results of the present study reveal that the quality of life of patients with facial prostheses is 'fair' or 'good'. Pain, activity, and appearance were considered the major issues affecting their quality of life.

This study was conducted using a validated UoW-QOL v4, as this was able to provide clinical information briefly and simply.¹⁴ The UoW-QOL has been translated into, and validated in, various languages by the Merseyside Regional Head and Neck Cancer Centre (2019). These include Brazilian,¹⁵ Spanish,¹⁶ Greek,¹⁷ Turkish,¹⁸ and many more. In

Table III: Percentage of patients choosing each domain and had significant problem on each UoW-QOL domain (n=64)

UoW-QOL	Patients choosing the domain, n (%)	Rank order	Patients with significant problems, n (%)
Pain	58 (90.6)	1	1 (1.6)
Activity	56 (87.5)	2	4 (6.3)
Appearance	50 (78.1)	3	9 (14.1)
Recreation	13 (20.3)	4	5 (7.8)
Anxiety	9 (14.1)	5	2 (3.1)
Mood	5 (7.8)	6	7 (11.0)
Swallowing	0 (0)	7	0 (0)
Chewing	0 (0)	7	0 (0)
Speech	0 (0)	7	0 (0)
Taste	0 (0)	7	0 (0)
Shoulder	0 (0)	7	0 (0)
Saliva	0 (0)	7	0 (0)

this study, the researchers used the translated and validated UoW-QOL questionnaire in Malay, which was sourced from the Merseyside Regional Head and Neck Cancer Centre official website.¹⁹ A self-administration mode similar to the study published by Kazi *et al.* (2008) was adopted since it avoids potential interview bias and is quick, simple, and convenient for patients to complete.²⁰

This study comprised 64 respondents, the majority of whom were male (64.1%). This was consistent with the findings of most studies of patients with a facial prosthesis because the incidences of head and neck cancers are around three times higher in males.²⁰ Moreover, about three-quarters (73%) of all road traffic accidents occur among young males under the age of 25. They are almost three times as likely to be involved in a road traffic collision compared to young females.²² The incidence of head and neck cancers in children was reported as having a male-to-female ratio of 1.78:1.²³ The current study reported patients had a mean age of 38.1 years, which differed slightly from the average, as head and neck cancers are known to have a higher incidence in individuals over 45 years old.²⁴ This was probably due to the inclusion criteria, which not only consisted of patients wearing facial prostheses due to head and neck cancers (53.1%) but also accidents (43.8%) and congenital conditions (3.1%).

Of the patients interviewed, 84.4% quoted their overall quality of life as 'fair' or 'good'. This result matches very closely to a study conducted by Kazi *et al.* (2008) in the United Kingdom, in which 43.7% of their respondents reported their quality of life as 'very good' and 28.1% reported it as 'good'.²⁰ The slight variation between these two studies in terms of the results obtained was probably because the United Kingdom was ranked second in a study comparing the healthcare systems of seven industrialised countries.²⁵

Pain (90.6%), activity (87.5%), and appearance (78.1%) were the issues that most affected patients, as reported in this series of interviews. The pain domain score is consistent with a study by Iriya P. *et al.* (2017), which reported pain as one of the most important issues experienced by patients.²⁶ This was because the head and neck region is highly susceptible to pain, due to extensive innervation and the proximity of anatomical structures.²⁷

The findings of the current study showed that the most significant problems encountered by the patients were appearance (14.1%), mood (11%) and recreation (7.8%). The finding for the recreation domain in this study is consistent with the study by Rogers *et al.* (2003), which reported patients with larger defects gave lower scores for activity, recreation, and physical function. On the other hand, the results for the mood and recreation domains are consistent with a study conducted by Dzebo S. *et al.* (2017), which pointed out that mood (53%), anxiety (43%), and recreation (39%) were the most significant problems encountered by patients.²⁸ These variations between the results obtained were possibly due to several associated factors such as gender, schooling, family income, and age.¹⁵

The UoW-QOL scale is brief and simple.²⁰ The researchers found that it can easily be used to assess patients with a facial prosthesis. It helps the clinician with useful data and can guide the clinician in decision-making based on patient feedback.²⁹ On the other hand, the UoW-QOL scale is based completely on the patient's perspective, which may cause inaccuracy in the results.²⁰ After conducting the study using the UoW-QOL questionnaire, the researchers would like to suggest improving the questionnaire by having an open-ended text that would allow patients to share their thoughts, and allow the researchers to recognise healthcare problems

that require attention but which may not have been identified beforehand.

CONCLUSION

The quality of life of patients with facial prostheses assessed using the UoW-QOL v4 scale was scored as 'fair' or 'good'. Pain, activity, and appearance were considered the major issues affecting their quality of life. To overcome the weaknesses of facial prostheses, proper instructions and sufficient information about the prostheses should be given to patients. Routine psychological consultations should also be offered to patients with facial prostheses.

ACKNOWLEDGEMENTS

The submission of this manuscript was supported by Universiti Sains Malaysia Incentive Bridging Grant 304.PPSG.6316453.

ETHICAL CONSIDERATIONS

The ethical clearance for this study was from the Human Research Ethics Committee of USM (JEPeM) (USM/JEPeM/19010035).

DECLARATION OF CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

REFERENCES

- 1: Aguiar L, Roberto Mozzini, A., Lersch, E., & De Conto, F. Palatal obturator: making an unconventional prosthesis - case report. *RFO UPF*. 2013;18(1):125-9.
- 2: Leonardi A, Buonaccorsi S, Pellacchia V, Moricca LM, Indrizzi E, Fini G. Maxillofacial prosthetic rehabilitation using extraoral implants. *J Craniofac Surg*. 2008;19(2):398-405.
- 3: Jeon B, Lee C, Kim M, Choi TH, Kim S, Kim S. Fabrication of three-dimensional scan-to-print

- ear model for microtia reconstruction. *J Surg Res*. 2016;206(2):490-7.
- 4: Horlock N, Vögelin E, Bradbury ET, Grobbelaar AO, Gault DT. Psychosocial outcome of patients after ear reconstruction: a retrospective study of 62 patients. *Ann Plast Surg*. 2005;54(5):517-24.
- 5: Rolski D, Kostrzewa-Janicka J, Zawadzki P, Życińska K, Mierzwińska-Nastalska E. The Management of Patients after Surgical Treatment of Maxillofacial Tumors. *Biomed Res Int*. 2016; 2016:4045329-.
- 6: Cole EBaCF. Children with hearing loss: Developing listening and talking, birth to six.: Plural Publishing; 2015.
- 7: Goiato MC, Pesqueira AA, Ramos da Silva C, Gennari Filho H, Micheline Dos Santos D. Patient satisfaction with maxillofacial prosthesis. Literature review. *Journal of plastic, reconstructive & aesthetic surgery: JPRAS*. 2009;62(2):175-80.
- 8: McKinstry RE. *Fundamentals of Facial Prosthetics*: ABI Professional Publications; 1995.
- 9: Ranabhatt R, Singh K, Siddharth R, Tripathi S, Arya D. Color matching in facial prosthetics: A systematic review. *J Indian Prosthodont Soc*. 2017;17(1):3-7.
- 10: Udagama A. Urethane-lined silicone facial prostheses. *J Prosthet Dent*. 1987;58(3):351-4.
- 11: Nemli SK, Aydin C, Yilmaz H, Bal BT, Arici YK. Quality of life of patients with implant-retained maxillofacial prostheses: a prospective and retrospective study. *T J Prosthet Dent*. 2013;109(1):44-52.
- 12: Deprich R, Naujoks C, Lind D, et al. Evaluation of the quality of life of patients with maxillofacial defects after prosthodontic therapy with obturator prostheses. *Int J Oral Maxillofac Surg*. 2011;40(1):71-9.
- 13: Becker C, Becker AM, Dahlem KKK, Offergeld C, Pfeiffer J. Aesthetic and functional outcomes in patients with a nasal prosthesis. *Int J Oral Maxillofac Surg*. 2017;46(11):1446-50.
- 14: Weymuller EA, Jr., Alsarraf R, Yueh B, Deleyiannis FW, Coltrera MD. Analysis of the performance characteristics of the University of Washington Quality of Life instrument and its modification (UW-QOL-R). *rch Otolaryngol Head Neck Surg*. 2001;127(5):489-93.
- 15: Andrade FPd, Biazevic MGH, Toporcov TN, Togni J, Carvalho MBd, Antunes JLF. Validade discriminante do questionário de qualidade de

- vida da Universidade de Washington no contexto brasileiro. *Revista Brasileira de Epidemiologia.* 2012; 15:781-9.
 - 16: Nazar G, Garmendia ML, Royer M, McDowell JA, Weymuller EA, Jr., Yueh B. Spanish validation of the University of Washington Quality of Life questionnaire for head and neck cancer patients. *Otolaryngol Head Neck Surg.: Official journal of American Academy of Otolaryngology-Head and Neck Surgery.* 2010;143(6):801-807, 7.e1-2.
 - 17: Linardoutsos G, Rapidis AD, Lowe D, Bramis I, Rogers SN. Development of the Greek version of the University of Washington quality of life questionnaire for patients with head and neck cancer. *J Craniomaxillofac Surg.: official publication of the European Association for Cranio-Maxillo-Facial Surgery.* 2014;42(5):601-7.
 - 18: Senkal HA, Hayran M, Karakaya E, Yueh B, Weymuller EA, Jr., Hoşal A. The validity and reliability of the Turkish version of the University of Washington Quality of Life Questionnaire for patients with head and neck cancer. *Am J Otolaryngol.* 2012;33(4):417-26.
 - 19: Merseyside Regional Head and Neck Cancer. UW-QOL v4 Translations [Internet]. 2019. Available from: <http://www.headandneckcancer.co.uk/professionals/quality-life/qol-questionnaires/uw-qol/uw-qol-v4-translations>.
 - 20: Kazi R, Johnson C, Prasad V, et al. Quality of life outcome measures following partial glossectomy: Assessment using the UW-QOL scale. *J Cancer Res Ther.* 2008;4(3):116-20.
 - 21: Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018;68(6):394-424.
 - 22: Organization. WH. 2020. Available from: <https://www.who.int/news-room/fact-sheets/detail/road-traffic-injuries>.
 - 23: Sengupta S, Pal R, Saha S, Bera SP, Pal I, Tuli IP. Spectrum of head and neck cancer in children. *J Indian Assoc Pediatr Surg.* 2009;14(4):200-3.
 - 24: Boyle P LB. World cancer report.: IARC Press: Lyon, France; 2008.
 - 25: Roe AM, Liberman A. A Comparative Analysis of the United Kingdom and the United States Health Care Systems. *Health Care Manag (Frederick).* 2007;26(3):190-212.
 - 26: Iriya PMO, Romaniszen LW, Fernandes TMF, Poleti ML. Health-related quality of life of patients with squamous cell carcinoma: a comparison according to tumor location. *Braz Oral Res.* 2017;31: e105.
 - 27: Chua KS, Reddy SK, Lee MC, Patt RB. Pain and loss of function in head and neck cancer survivors. *J Pain Symptom Manage.* 1999;18(3):193-202.
 - 28: Dzebo S, Mahmutovic J, Erkocevic H. Quality of Life of Patients with Oral Cavity Cancer. *Mater Sociomed.* 2017;29(1):30-4.
 - 29: Rogers SN, Lowe D, Brown JS, Vaughan ED. The University of Washington head and neck cancer measure as a predictor of outcome following primary surgery for oral cancer. *Head Neck.* 1999;21(5):394-401.
-