

EP_A097

SEVERE OSTEOPOROSIS WITH FRAGILITY FRACTURE REVEALING PRIMARY HYPERPARATHYROIDISM

Sarojini Devi Simanchalam¹, Poh Shean Wong¹, Nor Afidah Abdul Karim¹, Noor Lita Adam¹, Fauzi Azizan²

¹Department of Medicine and Endocrinology, Hospital Tuanku Jaafar

²Department of Medicine, Hospital Tuanku Ampuan Najihah

INTRODUCTION

Primary hyperparathyroidism (PHPT) is frequently asymptomatic or detected incidentally; however, delayed diagnosis may lead to severe skeletal complications. Early recognition is essential, as timely identification and management of parathyroid disease can prevent significant morbidity, although diagnosis may be challenging when clinical and imaging findings are inconclusive.

CASE

A 46-year-old female with severe bilateral hearing impairment presented to the orthopedic clinic with 3 years' history of bilateral knee pain and was found to have a right intertrochanteric femur fracture following minimal trauma. She was referred for evaluation of suspected secondary osteoporosis. She has had intermittent constipation and long-standing oligomenorrhea since menarche. There was no history of childhood fractures or use of medications affecting bone metabolism. Examination revealed bilateral knee bowing.

Initial evaluation considered metabolic bone disease, including Paget's disease; however, skeletal survey showed no features suggestive of Paget's disease or multiple myeloma.

Biochemical investigations demonstrated persistent hypercalcemia (2.65–3.1 mmol/L) with inappropriately elevated intact parathyroid hormone (peak 7.62 pmol/L), consistent with PHPT. Serum phosphate was low-normal. Concomitant vitamin D deficiency (25-OH vitamin D 34.46 nmol/L) improved following replacement. Bone mineral density confirmed severe osteoporosis (lumbar spine T-score -5, z-score -4.1); (forearm -6.7, z-score -6.1) reflecting prolonged untreated disease.

Neck ultrasound demonstrated a mixed solid-cystic lesion posterior to the right thyroid lobe, suggestive of a parathyroid adenoma. The TC-99 m Sestamibi scan showed no definite focal uptake. However, subsequent SPECT-CT revealed focal tracer uptake at the posterior right thyroid gland, consistent with a hyperfunctioning parathyroid gland. Parathyroidectomy was done. Postoperatively, the calcium level normalized.

CONCLUSION

Severe osteoporosis and fragility fracture occur, reflecting prolonged exposure to excess parathyroid hormone and significant skeletal morbidity. Early biochemical evaluation in unexplained severe osteoporosis is essential, as timely diagnosis and definitive management of parathyroid disease are critical to halt ongoing bone loss and prevent irreversible complications.

EP_A098

A CHALLENGING CASE OF PARATHYROID CARCINOMA IN AN ADOLESCENT

S. Muhammad Imran¹ and Tong Chin Voon¹

¹Hospital Putrajaya

INTRODUCTION

Parathyroid carcinoma (PC) is an exceedingly rare malignancy. Definitive surgical management with en bloc resection is crucial for cure, but the post-operative course can be complicated by profound metabolic derangements, most notably hungry bone syndrome (HBS). We report a case of PC in a 15-year-old female to highlight the challenges in perioperative management.

CASE

A 15-year-old female presented with a painless, palpable neck mass. There is no family history of note. Investigations revealed severe primary hyperparathyroidism (PHPT) with a corrected calcium of 3.4 mmol/L and an elevated intact parathyroid hormone level of 118.6 pg/mL. Alkaline phosphatase was markedly elevated at 1,168 U/L. Ultrasound, computed tomography neck, and a sestamibi scan identified a large, lobulated 4.7-cm mass posterior to the right thyroid lobe, suspicious of malignancy. Bone mineral density of the forearm was severely diminished, with a Z-score of -6.6. The patient underwent right hemithyroidectomy and parathyroidectomy with intraoperative neural monitoring. Intraoperative parathyroid hormone levels dropped from a pre-excision level of 120.1–16.5 pg/mL 5 minutes post-excision, confirming complete resection of the hyperfunctioning tissue. Histopathological examination confirmed the diagnosis of PC, demonstrating lymphovascular invasion and clear resection margins. The Ki-67 proliferation index was 5%. Post-operatively, the patient developed hypocalcemia, with corrected calcium dropping to a nadir of 1.95 mmol/L. This was managed with intensive calcium and activated vitamin D supplementation. At 10 months post-surgery, the patient continues to require supplementation for persistent hypocalcemia. Surveillance ultrasound at 3 months showed no evidence of recurrence, and she is planned for ongoing annual monitoring.