

ORIGINAL ARTICLE

IMPACTS OF DIETARY SUPPLEMENTS AND NUTRIENT-RICH FOOD FOR PREGNANT WOMEN ON BIRTH WEIGHT IN SUGH EL-CHMIS /ALKHOMS - LIBYA

Alabed Ali A. Alabed¹, Eman Ali Aljrbi², Abdullah Y. Al-Mahdi^{*3}, Rasheed Abdulsalam⁴, Anis Rageh Al-Maleki⁵ and Hasanain Faisal Ghazi⁶

¹Department of Community Medicine, Faculty of Medicine, Lincoln University College, 47301 Petaling Jaya, Selangor, Malaysia

²Department of Biotechnology, Faculty of Science, Lincoln University College, 47301, Petaling Jaya, Selangor, Malaysia

³Department of Microbiology, Faculty of Medicine, Lincoln University College, 47301, Petaling Jaya, Selangor, Malaysia

⁴Faculty of Dentistry, Lincoln University College, 47301, Petaling Jaya, Selangor, Malaysia

⁵Medical Microbiology Department, Faculty of Medicine, University of Malaya, Kuala Lumpur, 50603, Malaysia

⁶College of Nursing, Al-Bayan University, Baghdad, Iraq

*Corresponding author: Abdullah Y. Al-Mahdi

Email: dr.microbiology2019@gmail.com

ABSTRACT

This study aims to determine the prevalence of low birth weight in Sugh El-Chmis /Alkhoms -Libya and the associated factors with the low birth weight of the baby. A cross-sectional study was performed in Libya, in 2015 using a pre-tested self-administered questionnaire on a stratified sample of 408 pregnant women. Descriptive, bivariate and multivariate analyses were carried out for statistical analysis. Out of a total of 408 respondents, the prevalence of low birth-weight baby was 23.5%. The most of participants were aged between 19-26 years old (49%). Dietary supplements (folic acid, iron, omega and vitamin A, D, E, S, M), natural fruits (apple, grape, watermelon, plum, figs, strawberry and peaches), all milk and dairy products (milk, cheese and yogurt), meat, fish, egg, legumes were significantly associated with baby weight ($P < 0.05$) using bivariate analysis. Furthermore, other factors such as culture and lifestyle and popular diets such as breakfast time, additional meal, soft drink, stimulants, barley wheat brown bread and dates also closely associated with the low-birth weight of new-borns ($P < 0.05$). The intake of dietary supplements and nutrient-rich food for pregnant women influenced on the birth weight in Sugh El-Chmis /Alkhoms-Libya. Therefore, awareness regarding intake of dietary supplements and nutrient-rich food is highly recommended.

Keywords: Dietary supplementary, Nutrient-rich food, Libya, Low birth weight.

INTRODUCTION

Globally, more than 20 million infants representing 15.5% of all births, are born with low birth weight, 95.6% of them in developing countries. The level of low birth weight in developing countries (16.5%) is more than double the level in developed regions (7%). Half of all low-birth-weight babies are born in South-central Asia, where more than a quarter (27%) of all infants weighs less than 2,500 g at births¹.

Maternal nutrition and health are considered the most important regulator of human fetal growth. Improved maternal nutrition has been associated with increased fetal growth and a reduction in adverse birth outcomes in developing countries and populations with nutrient deficiencies². However, if women are not well nourished, they are more likely to give birth to weak babies resulting in a high infant mortality rate. At birth, fetal weight is accepted as a single parameter that is directly related to the health and nutrition of the mother. On the other hand, it is an important determinant of the chances of the newborn to survive and experience healthy growth and development. This is because low

birth weight is directly related to both immediate, long-term and very long-term development and well-being³.

World Health Organization (WHO) has defined low birth weight (LBW) as weight at birth of less than 2500 g⁴. This definition of LBW includes in its total a subgroup of infants who have suffered varying degrees of nutritional deprivation in utero. The latter subgroup, victims of intrauterine growth retardation (IUGR), constitutes the group at the highest degree of risk of both short-term and long-term complications. Fetal growth and birth weight are influenced by a variety of factors, racial, social and economic among others, as well as specific medical conditions that may be present or that may develop during pregnancy⁵.

A few investigations have demonstrated a relationship between anthropometric indicators and pregnancy outcomes. Both inadequate and unreasonable gestational weight gain is strongly connected with maternal-fetal complexities, for example, gestational diabetes, hypertensive pregnancy issue (HPD), macrosomia, and low birth weight⁶. Contrasted with ordinary weight, maternal overweight is identified with a higher

danger of cesarean deliveries and a higher rate of aesthetic and postoperative complications in these deliveries. Low Apgar scores, macrosomia, and neural tube defects are more frequent in newborn children of fat mothers than in babies of ordinary weight moms⁷. Inadequacies in micronutrients may influence baby birth weight and survival, and poor vitamins. An intake builds the danger of maternal night visual deficiency⁸. Nutritional status in pregnancy is a basic issue that can't be neglected while considering the wellbeing of a nation, particularly the developing. A woman's dietary status has essential ramifications for her well-being and the well-being of her kids. Malnutrition in women brings about lessened efficiency, an expanded weakness to contaminations, moderate recuperation from sickness, and elevated dangers of unfriendly pregnancy results. For instance, a woman who has poor nutritious status, as shown by a low body mass index (BMI), short stature, or other micronutrient inadequacies, has more danger of deterred work, of having an infant with low birth weight, of delivering lower quality breast milk, of passing on from baby blues discharge, and of contracting maladies alongside her infant⁹. The brilliant interim for mediation stretches out from pregnancy to two years old. On the off chance that undernutrition isn't tended to amid these critical times of life, it might cause irreversible harm for future improvement towards adulthood, hence expanding the danger of young ladies getting to be malnourished moms, who at that point have a low-birth-weight infant.

The percentage and number of low-birth-weight infants by UNICEF was published in 2000, showed that 15% was estimated of low birth weight in the Middle East and North Africa with 60% of birth not weighed. The same result was reported for Western and Central Africa and a little less in Eastern and Southern Africa by 14%. In Libya, it was almost half by 7% of low-birth-weight infants for 1000¹. Moreover, stressful life events experienced by pregnant women may lead to adverse obstetric outcomes. A study in Benghazi compared the rates of preterm, low-birth-weight and cesarean-section births at Al-Jamhouria hospital in the months before and during the armed conflict in Libya in 2011. Compared with the preceding months there was a significant rise during the conflict in the rate of deliveries involving preterm (3.6% versus 2.5%) and low-birth-weight (10.1% versus 8.5%) infants and cesarean sections (26.9% versus 25.3%).

According to the most recent WHO data from 2018, low birth weight fatalities in Libya totaled 389, accounting for 1.24 percent of all deaths. Libya ranks at the 109 positions in the world with an age-adjusted death rate of 5.48 per 100,000 people. Between 1995 and 2014, the prevalence of underweight, weight for age, male (percentage of children under the age of 5) in Libya increased

significantly from 4.6 to 13%, climbing at an annual rate of 106.35 percent. Therefore, this study aims to determine the prevalence of low birth weight in Sugh El-Chmis /Alkhoms -Libya and the associated factors with the birth weight of the baby.

MATERIALS & METHODS

Study design and sampling

A cross-sectional study was conducted among pregnant women who were in their first, second or third trimester visiting Women and Children government hospital in Sugh El-Chmis, Libya. The pregnant women who used to take dietary supplements and natural food during their pregnancy were chosen to participate in this study. People with the ability to read and write the questionnaire considered eligible for this study. Participating in this study was voluntary and random. A total of 408 questionnaires were collected. All respondents were briefed about the study at the beginning of the questionnaire as a short introduction. The questionnaires were given to pregnant women after obtained verbal consent from them. Only those who fulfilled the inclusion and exclusion criteria and consented were included in this study. The data excel file was obtained by the authors after it was taken from the Google form.

Measurement tool

A primary questionnaire was developed after a comprehensive literature review and validated by a panel of experts consisting of an epidemiologist, nutrient specialist, public health specialist and biostatistician to be used for pregnant women who visiting Women and Children Government Hospital at Sugh El-Chmis, Libya. A pre-test was performed at ALMGREF health center, Libya to make sure the questions are clear and easy to understand. After the approval of the pre-testing, the survey was performed via a questionnaire. The questions were prepared in the Arabic language and back-translated into the English language. The questionnaires were divided into five sections including economic and social situation; dietary supplements; natural food; culture and lifestyle and popular diets; and birth weight and maternal and child health status. Most of the items were designed to be closed-ended questions by answering yes or no. The questionnaire was designed to be anonymous, and informed consent was obtained from every respondent as it was sent along with the survey. The data were kept confidential and the results did not identify the respondents personally. Participating in the research was voluntary and anonymous.

Data analysis

Data analysis was performed using Statistical Package for Social Sciences (SPSS) version 22

(Chicago, IL, USA). Continuous variables were expressed as mean $\pm$ standard deviation (SD) while categorical variables were presented as frequency and percentage. Descriptive statistics were carried out to determine the socio-demographic factors and baby weight. Inferential statistics (chi-square test) was used to assess the association between socio-demographic factors and baby weight. Then, multivariate analysis was performed to identify the most important factors associated with baby weight. Multiple logistic regression was used for this analysis. The significance level was set at ($P \leq 0.05$) for all statistical procedures.

RESULTS

Socio-Demographic Data

Out of total of 408 participants in this study, the prevalence of low birth was 23.5%. The most of respondents were aged between 19-26 years old (49%), 112 (27.5%) aged between 27-36 years old, 72 (17.6%) aged more than 36 years old and 24 (5.9%) aged less than 19 years old. Women with less than 5 years of marriage were the majority by 208 (51.0%) compared to other groups which represent 40 (9.8%) for 5-10 years marriage, 72 (17.6%) for 10-15 years marriage and 88 (21.6%) for more than 15 years marriage. The majority of the respondents were university-educated with 232 (56.9%) while intermediate; and read and write accounted for 144 (35.3%) and 32 (7.8%) respectively. It was found that most of women 256 (62.7%) were from family with middle income 501-1000 LYD compared to 112 (27.5%) with less than 500 LYD, 32 (7.8%) with 1011-1500 LYD and 8 (2.0%) with more than 1500 LYD. The participants were mostly housewives by 200 (49.0%) followed by 16 (3.9%) teachers, 40 (9.8%) employees and 8 (2.0%) craftswomen. Finally, 248 (60.8%) women had 2-5 children compared to 88 (21.6%) one child and 72 (17.6%) 6-9 children. The summary of the characteristics is shown in Table 1.

Bivariate Analysis Related to Dietary Supplements

Chi-square test was used to determine the association between baby weight and all the categorical variables such as: dietary supplements, natural food, meat, eggs and legumes, culture and lifestyle and popular diets and maternal and child health status. It was found that dietary supplements (folic acid, iron, omega and vitamin A, D, E, S, M) were significantly associated with baby weight ($P < 0.05$). Only calcium was not statistically significant with baby

weight ($P > 0.05$). For vegetables, it was demonstrated that tomato, beef and cabbage were significantly associated with baby weight ($P < 0.05$) except lettuce, cucumber and carrots.

For natural fruits, it was revealed that apple, grape, watermelon, plum, figs, strawberry and peaches were significantly associated with baby weight ($P < 0.05$). On the other hand, banana, apricot, tin pear, orange and kiwi were not statistically significant with baby weight ($P > 0.05$). Interestingly, it was showed that all the dairy products (milk, cheese and yogurt), red meat, white meat, fish, eggs and all kind of legumes were significantly associated with baby weight ($P < 0.05$). Furthermore, it was demonstrated that all factors for culture and lifestyle and popular diets such as breakfast time, additional meal, soft drink, stimulants, barley wheat brown bread and dates were too significantly associated with baby weight ($P < 0.05$). The analysis indicated that newborn health status, mother health, delivery date and breast feeding were statistically associated with baby weight ($P < 0.001$) but only delivery status factor was not significant with baby weight ($P > 0.05$).

Multivariate analysis

Multiple logistic regression analysis was performed to estimate the predictive risk factors of baby weight when other confounders were controlled or adjusted so that they could be calculated. All the variables with a total sample size of 408, which were statistically significant in the bivariate analysis, were included in the multiple logistic regression analysis. The dependent variable that measures the baby weight was YES, which was coded as (1) for normal weight, while NO was coded as (0) for not normal.

As shown in Table 3, logistic regression analysis revealed that the factors that were significantly associated with baby weight after controlling the confounders ($P < 0.05$) by using stepwise forward LR logistic regression method. The results show that the dates and breastfeeding ($P < 0.001$) made a significant contribution to prediction, whereas other variables were not. The most important factor responsible for the estimation of the baby weight was identified by using the Wald criterion, which gave the 'importance' of the contribution of each variable in the model. The Wald test demonstrated that the dates was the most important predictor for baby weight estimation.

Table 1: Descriptive analysis related to demographical factors

Variables	Frequency	Percent
Age		
Less than 19	24	5.9
19-26	200	49.0
27-36	112	27.5
More than 36	72	17.6
Duration of marriage		
Less than 5	208	51.0
5-10	40	9.8
10-15	72	17.6
More than 15	88	21.6
Educational level		
Intermediate educ.	144	35.3
University educ.	232	56.9
Read and write	32	7.8
Family income		
Less than 500 LYD	112	27.5
501-1000	256	62.7
1001-1500	32	7.8
More than 1500 LYD	8	2.0
Occupation		
Housewife	200	49.0
Employee	40	9.8
Teacher	160	39.2
Craftswomen	8	2.0
No. of children		
One child	88	21.6
2-5	248	60.8
6-9	72	17.6
TOTAL	408	100.0

Table 2a: Bivariate analysis for most important factors related to baby low birth weight

Variables	X ² test	P value
<u>Dietary supplementary</u>		
Folic acid	17.018	0.002*
Calcium	6.613	0.307
Iron	28.714	0.000*
Omega	23.099	0.000*
Vit. A, D, E, S, M	12.669	0.013*
<u>Vegetables</u>		
Lettuce	2.931	0.401
Tomatoes	15.551	0.019*
Cucumber	6.221	0.101
Beet	50.158	0.000*
Carrots	4.670	0.198
Cabbage	49.244	0.000*
<u>Fruits</u>		
Apple	54.697	0.000*
Banana	4.222	0.238
Grape	17.468	0.001*
Watermelon	10.440	0.015*
Apricot	3.193	0.363
Tin pear	7.553	0.056*
Plum	11.342	0.010*
Figs	13.293	0.004*
Orange	7.172	0.067
Kiwi	6.417	0.093
Strawberry	32.579	0.000*
Peaches	11.613	0.009*

Table 2b: Bivariate analysis for most important factors related to baby low birth weight

Variables	X ² test	P value
<u>Milk and dairy products</u>		
Milk	28.071	0.000*
Cheese	24.360	0.000*
Yogurt	66.771	0.000*
<u>Meat, fish, eggs and legumes</u>		
Red meat	64.099	0.000*
White meat	58.820	0.000*
Fish	8.171	0.029*
Eggs	22.789	0.003*
All kind of legumes	22.280	0.001*
<u>Culture, lifestyle and popular diets</u>		
Breakfast time	40.917	0.000*
Additional meal	66.413	0.000*
Soft drink	30.775	0.000*
Stimulants	26.673	0.000*
Meal by barley wheat brown bread	69.424	0.000*
Dates	89.747	0.000*
<u>Maternal and child health status</u>		
Newborn health status	8.938	0.011*
Mother health	23.050	0.000*
Delivery date	16.299	0.001*
Delivery status	0.090	0.887
Breast feeding	37.430	0.000*

*p<0.05

Table 3: Multivariate Analysis for Most Important Factors Related to Baby Low Birth Weight

Variables	B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for EXP(B)	
							Lower	Upper
Dates			93.157	3	0.000*			
Daily	3.014	0.417	52.298	1	0.000*	20.363	8.997	46.086
2-3 times a week	1.275	0.237	28.923	1	0.000*	3.579	2.249	5.695
3-4 weeks	4.152	0.553	56.410	1	0.000*	63.536	21.503	187.727
Breastfeeding			50.357	2	0.000*			
Normal	-0.440	0.255	2.975	1	0.085	0.644	0.391	10.062
Non -normal	-3.673	0.518	50.353	1	0.000*	0.025	0.009	0.070

*p<0.05

DISCUSSION

To the best of our knowledge, this is the first time that a study demonstrates the prevalence of low birth weight and their associated factors in Sugh El-Chmis, Libya. The findings can be considered the first step in providing data on low birth weight and will help in creating awareness among Libya pregnant women, to provide insight in designing future action steps to bridge the gaps as an effort against low birth weight. In this study, we found that there was a relationship between taking protein day by day and newborn childbirth weight ($p < 0.05$). This outcome was coordinated with the investigation of Whitney et al. (2007) study which demonstrated that protein is important for the fast fetal development, the extension of the uterus, mammary organs, and placenta,

increments in maternal blood volume, development of amniotic liquid, and capacity save for work, conveyance, and lactation¹⁰. Many shreds of evidence have shown that macronutrients (protein) supplementation were associated with an increase in maternal weight and mean birth weight and a decrease in the number of low birth weight babies. According to the World Health Organization (1995), an additional 100 kcal/day throughout the pregnancy can increase birth weight by about 100 g and the risk of IUGR would be halved for undernourished women¹¹.

Poor iron status may affect immune function and thus increase host susceptibility to "genital tract infection" and also increase oxidative stress of the placenta. Low hemoglobin status causes chronic

hypoxia¹². Supplementation of 60 mg of ferrous iron and 0.4 mg of folic acid /day significantly increased the birth weight. To obtain a maximum effect on birth weight iron supplementation should be started in early pregnancy¹³. Iron and folic acid deficiency in early pregnancy are associated with neurological defects. In this study, it proved that iron has an association with baby weight ($p < 0.05$) and supported the results from Muthayya et al. (2006) study¹³. Prevalence of anemia among pregnant women in Libya is comparable to that of a severe public health concern, indicating that anemia in pregnant women is a public health issue in Libya. According to Elzahaf and Omar (2016) study, it has been reported that anaemia was more common in pregnant women aged 15-20 to 36-40 years old and with only a primary school education. Anemia was found in 54.6 percent of pregnant women, with 44.5 percent having mild anemia. The third trimester of pregnancy has a higher prevalence of anemia (59.6 percent). Thus, awareness regarding regular intake of iron and folic acid tablet was highly recommended in their study. Furthermore, vitamin A is fundamental for development, and the separation of most cells and tissues. It is essential for sound fetal advancement. Pregnant women are by and large encouraged to expand their vitamin intake by 10- 40%¹⁴.

Calcium is important because it aids to strengthen the baby's rapidly-developing bones and teeth, and boosts muscle and heart. It plays a vital role in nerve development too. Moreover, it is still essential for teeth and bones of a pregnant woman. If the pregnant woman doesn't get enough calcium in their diet, the body will take what the baby needs. It is especially the case during the third semester when bone development peaks at 250-250 mg transferred from the pregnant woman and the baby every day¹⁵. A woman who is not getting enough calcium during her pregnancy makes her more susceptible to osteoporosis. Unlike our study demonstrated that calcium was not statistically significant ($p = 0.307$). However, concerning the dietary supplements, this investigation shows that there was a factual huge positive relationship between taking supplements (eg. omega and vitamins) and new-born childbirth weight. This outcome was bolstered by Lassi et al. (2006) study¹⁶. They found that numerous micronutrient supplementations brought about a measurably diminished number of low-birth-weight babies, small for gestational age (SGA) babies and maternal pallor.

Omega-3 fatty acids are essential and can only be obtained from the diet. Nevertheless, a great many people, including pregnant women, are not getting sufficiently omega-3 from their eating regimen. Omega-3 unsaturated fats are basic amid pregnancy, particularly the long-chain omega-3 unsaturated fats DHA and EPA. These are

found in high sums in fish and help assemble the cerebrum and eyes of the baby. Pregnant women are encouraged to confine their fish intake for twice per week. Due to the mercury and different contaminants found in greasy fish, this has made a few women maintain a strategic distance from fish by and large, in this way restricting the intake of fundamental omega-3 unsaturated fats. Nonetheless, studies have demonstrated that pregnant women who eat 2-3 dinners of greasy fish every week accomplish the suggested intake of omega-3 and increment their blood levels of EPA and DHA. Salmon is likewise one of not very many common wellsprings of vitamin D, which is regularly ailing in the eating routine. It is imperative for some procedures in the body, including bone wellbeing and resistant capacity¹⁷. In our study, it revealed that omega was significantly associated with low birth weight ($p = 0.000$). Based on Albakoush and Azab (2019) study, it has been demonstrated that pregnant women in Western-Libya were at an increased risk of vitamin D insufficiency. Pregnant women's vitamin D levels were found to be much lower than non-pregnant women's. Vitamin D levels were below 20ng/ml in 84.8 percent of the study population, while calcium levels were below 8.4 mg/dl in 46.8%. Vitamin D levels were less than 20ng/ml in 95.2 percent, 91.2 percent, and 66.7 percent of pregnant women in the first, second, and third trimesters, respectively. Calcium levels were less than 8.4 mg/dl in 61.9 percent, 35.3 percent, and 50 percent of pregnant women in the first, second, and third trimesters, respectively. The study focused on maternal nutrition, particularly optimal vitamin D and calcium intake, which may prepare the way for future bone health concerns such as osteoporosis prevention in the long term. Moreover, vegetables and fruits are one of the most important nutrients that should be taken by pregnant women as suggested by the World Health Organization (WHO). Fresh vegetables and fruits contain lots of essential vitamins and nutrients and are a good source of fiber too. This not only helps the fetal development but also keep the pregnant woman body in the best condition possible. In this study, the importance of these nutrients was very clear through our results which demonstrated that most of them were statistically significant and have an association with baby weight ($p < 0.05$).

Red and white meat is a great source of protein, zinc and iron. These are essential nutrients that every pregnant woman should consume in adequate amounts during pregnancy. Expanding fish was more marginally among the mothers who brought forth LBW new-born children. This implies that there is a negative connection between the utilization of fish amid pregnancy and baby birth weight. This was due to the defilement with toxins and dangerous metals as mercury which might be kept in angle. This toxic compound may gather in the circulation system after some time

and could harm the new-born child developing cerebrum and sensory system. This is conflicting with Thorsdottir et al. (2004) study which discovered the baby of women in the most reduced amounts of fish utilization weighed less, were shorter and had a littler head outline during childbirth than those of women expending higher measures of fish¹⁸. Legumes are brilliant plant-based sources of fiber, protein, press, folate (B9) and calcium, all of which the body needs a greater amount of amid pregnancy. Folate is one of the B-vitamins (B9). It is imperative for the wellbeing of the mother and baby, particularly amid the primary trimester¹⁹. Eggs are rich in the mineral choline and Omega 3 fatty acids. They are important for baby's brain development. It helps to prevent neural tube defects *in vitro* as well.

In this study, culture, lifestyle and popular diets (soft drink, stimulants, dates and barley, wheat brown bread) significantly associated with low birth weight. According to Abubakari and Jahn (2016) study, skipping breakfast affects fetal development. Soft drinks have carbon dioxide bubbles²⁰. Carbonic acids will cause severe heartburn and indigestion, increasing the acidic levels in a pregnant lady. Barley wheat brown bread is a good source of the B vitamins (including B1, B2, folic acid, and niacin) which is crucial for baby development and growth of the placenta. Based on Karen Miles (2020) study, it has been demonstrated that the recommended amount of fiber during pregnancy is 28 g a day, which helps prevent constipation and haemorrhoids²¹.

Finally, dates are guaranteed as 'heart solid nourishments' by the American Heart Association because of their low cholesterol and immersed fat substance. They contain supplements that are fundamental to the baby's sound development and advancement. They have many advantages, for example, insurance of the liver, help vitality, enhances weight, support by insusceptible framework, enhances hemoglobin, enhances vision, cures looseness of the bowels and numerous others²². In this study, date one the most important and predictors for baby weight as it was statistically significant in the bivariate analysis and multivariate analysis as well along together with breastfeeding ($p < 0.05$). According to the World Health Organization (2020), breastfeeding has many health benefits for both the mother and baby²³. Breast milk contains all the nutrients a baby needs in the first six months of life. It protects against diarrhoea, allergies, sickness, obesity and infections.

CONCLUSION

The present findings highlight concerns regarding the intake of dietary supplements (folic acid, iron, omega and vitamin A, D, E, S, M)), natural fruits (apple, grape, watermelon, plum, figs, strawberry and peaches), milk and dairy products (milk,

cheese and yogurt), meat, fish, egg, legumes, influence the low birth weight of new-borns in Sugh El-Chmis /Alkhoms-Libya. Furthermore, other factors such as culture and lifestyle and popular diets such as breakfast time, additional meal, soft drink, stimulants, barley wheat brown bread and dates also causes the low-birth weight of new-borns. These findings will aid to design effective interventions to overcome the low birth weight and create a more robust awareness among pregnant women.

Competing Interests

The authors have declared that no competing interest exists.

REFERENCES

1. United Nations Children's Fund and World Health Organization, Low Birthweight: Country, regional and global estimates. UNICEF, New York, 2004.
2. Black RE, Victora CG, Walker SP, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The lancet*. 2013; 382(9890):427-51.
3. Agarwal K, Agarwal A, Agrawal VK, et al. Prevalence and determinants of " low birth weight" among institutional deliveries. *Annals of Nigerian Medicine*. 2011; 5(2):48.
4. WHO. Global nutrition targets 2025: low birth weight policy brief Geneva. World Health Organization; 2014. Available from: https://www.who.int/nutrition/publications/globaltargets2025_policybrief_lb w/en/.
5. Hoyme HE, May PA, Kalberg WO, et al. A practical clinical approach to diagnosis of fetal alcohol spectrum disorders: clarification of the 1996 institute of medicine criteria. *Pediatrics*. 2005; 115(1):39-47.
6. Geraci M. Estimation of regression quantiles in complex surveys with data missing at random: An application to birthweight determinants. *Statistical methods in medical research*. 2016; 25(4):1393-421.
7. Gunatilake RP, Perlow JH. Obesity and pregnancy: clinical management of the obese gravida. *American journal of obstetrics and gynecology*. 2011; 204(2):106-19.

8. World Health Organization. Vitamin A supplementation in postpartum women. 2009. Available from: https://www.who.int/elena/titles/vitamina_postpartum/en/
9. Imdad A, Lassi Z, Salaam R, et al. Prenatal nutrition and nutrition in pregnancy: Effects on long-term growth and development. In *Early Nutrition and Long-Term Health 2017* Jan 1 (pp. 3-24). Woodhead Publishing.
10. Whitney E, Rolfes SR. 2002. *Understanding Nutrition*. Ninth Edition Belmont: Wadsworth Com.
11. World Health Organization, Maternal anthropometry and pregnancy outcomes. *Bull. World Health Org.* 1995; 73(suppl):1-98.
12. Murphy JD, Aronovitz MJ, Reid LM. Effects of chronic in utero hypoxia on the pulmonary vasculature of the newborn guinea pig. *Pediatric research*. 1986; 20(4):292-5.
13. Muthayya S, Kurpad AV, Duggan CP, et al. Low maternal vitamin B 12 status is associated with intrauterine growth retardation in urban South Indians. *European journal of clinical nutrition*. 2006; 60(6):791-801.
14. Beaton GH. Effectiveness of vitamin A supplementation in the control of young child morbidity and mortality in developing countries. *United Nations Subcommittee on Nutrition (SCN) News*. 1993; 9:17-23.
15. Karla Walsh. How Much Calcium Do You Need During Pregnancy? (2020). Available from: <https://www.whattoexpect.com/pregnancy/diet/calcium-during-pregnancy/#:~:text=Calcium%20helps%20strengthen%20your%20baby's,take%20what%20your%20baby%20needs.>
16. Lassi ZS, Salam RA, Haider BA, et al. Folic acid supplementation during pregnancy for maternal health and pregnancy outcomes. *Cochrane Database of Systematic Reviews*. 2013(3).
17. Calder PC. Omega-3 polyunsaturated fatty acids and inflammatory processes: nutrition or pharmacology? *British journal of clinical pharmacology*. 2013; 75(3):645-62.
18. Thorsdottir I, Birgisdottir BE, Halldorsdottir S, et al. Association of fish and fish liver oil intake in pregnancy with infant size at birth among women of normal weight before pregnancy in a fishing community. *American journal of epidemiology*. 2004; 160(5):460-5.
19. Hjertholm KG, Iversen PO, Holmboe-Ottesen G, et al. Maternal dietary intake during pregnancy and its association to birth size in rural Malawi: A cross-sectional study. *Maternal & child nutrition*. 2018; 14(1):e12433.
20. Abubakari A, Jahn A. Maternal dietary patterns and practices and birth weight in Northern Ghana. *PLoS One*. 2016; 11(9):e0162285.
21. Karen Miles. Grains in your pregnancy diet. (2020). Available from: https://www.babycenter.com/0_grains-in-your-pregnancy-diet_1497.bc
22. Jadidi MY, Sang SJ, Lari H. The effect of date fruit consumption on spontaneous labor. *Journal of Pizhūhish dar dīn va salāmat*. 2016; 1(3):4-10.
23. World Health Organization 2020. Available from: https://www.who.int/elena/titles/exclusive_breastfeeding/en/